

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 347 488 B1**

(12)

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: 01.06.94 (51) Int. Cl.⁵: **C07D 207/42, C07D 207/34, C07D 405/04, C07C 255/34, C07C 215/28, A01N 43/36**
- (21) Application number: **88112810.2**
- (22) Date of filing: **05.08.88**

(54) **Arylpyrrole insecticidal acaricidal and nematocidal agents and method for the preparation thereof.**

- (30) Priority: **23.06.88 US 208841**
- (43) Date of publication of application:
27.12.89 Bulletin 89/52
- (45) Publication of the grant of the patent:
01.06.94 Bulletin 94/22
- (94) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE
- (56) References cited:

EP-A- 0 080 051	EP-A- 0 130 149
EP-A- 0 182 737	EP-A- 0 206 999
GB-A- 2 111 985	US-A- 3 864 491

- (73) Proprietor: **AMERICAN CYANAMID COMPANY**
One Cyanamid Plaza
Wayne, NJ 07470-8426(US)
- (72) Inventor: **Brown, Dale Gordon**
361 Stonybrook Road
Hunterdon, NJ(US)
Inventor: **Siddens, Jack Kenneth**
36B Berrien Avenue
Princeton Junction, NJ(US)
Inventor: **Diehl, Robert Eugene**
17 Clementon Way
Lawrenceville, NJ(US)
Inventor: **Wright, Donald Perry, Jr.**
RR No. 1, Box 294 B1,
Woosamonsa Road
Pennington, NJ(US)
- (74) Representative: **Wächtershäuser, Günter, Dr.**
Tal 29
D-80331 München (DE)


Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 347 488 B1

Description


The present invention is directed to certain novel arylpyrrole compounds that are highly effective insecticidal, acaricidal and nematocidal agents useful for the control of insect, acarid and nematode pests and for protecting agronomic crops, both growing and harvested, against the ravages of said pests. The present invention is also directed to methods for preparing the arylpyrrole compounds.

The novel arylpyrrole compounds of the present invention have the structural formula illustrated as formula I:



wherein X is F, Cl, Br, I, or CF₃; Y is F, Cl, Br, I, CF₃ or CN; W is CN or NO₂ and A is H; C₁-C₄ alkyl optionally substituted with from one to three halogen atoms, one hydroxy, one C₁-C₄ alkoxy or one C₁-C₄ alkylthio, one phenyl optionally substituted with C₁-C₃ alkyl or C₁-C₃ alkoxy or with one to three halogen atoms, one phenoxy optionally substituted with one to three halogen atoms or one benzyloxy optionally substituted with one halogen substituent; C₁-C₄ carbalkoxymethyl; C₃-C₄ alkenyl optionally substituted with from one to three halogen atoms; cyano; C₃-C₄ alkynyl optionally substituted with one halogen atom; di-(C₁-C₄ alkyl) aminocarbonyl; or C₄-C₆ cycloalkylaminocarbonyl; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:


-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCF₃, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2.


The term C₄-C₆ cycloalkylamino carbonyl means a C₄ to C₆ cycloalkylamino group attached directly to the carbonyl group through the nitrogen atom.

A preferred group of novel arylpyrroles of the present invention are illustrated by formula II:




wherein A, L, M, R, W, X and Y are as described above.

Another preferred group of novel arylpyrroles of this invention are represented by formula III:




10 wherein A, L, M, R, W, X and Y are as described above.

Another group of preferred arylpyrroles of the invention are depicted by formula IV:




25 wherein A, L, M, R, W, X and Y are as described above.

Yet another group of preferred arylpyrroles of this invention are delineated by formula V:



40 wherein A, L, M, R, W, X and Y are as described above; and still other preferred arylpyrroles of the invention are depicted by formulas VI and VII:



55 wherein A, L, M, R, W, X and Y are as described above.

Preferred formula I arylpyrroles of the invention are those in which A is hydrogen or C₁-C₄ alkoxymethyl; W is CN or NO₂; L is hydrogen or F; X and Y are each Cl, Br or CF₃; M is H, F, Cl or Br; and R is F, Cl, Br, CF₃ or OCF₃.

Preferred formula II compounds which are especially effective as insecticidal, acaricidal and/or nematocidal agents are those in which A is hydrogen or C₁-C₄ alkoxymethyl; L is hydrogen; M is hydrogen, F, Cl or Br; R is F, Cl, Br, CF₃ or OCF₃; W is CN and X and Y are each independently Cl, Br or CF₃.

Other formula II compounds that are highly effective as insecticidal, acaricidal and/or nematocidal agents are those in which A is hydrogen or C₁-C₄ alkoxymethyl; L is hydrogen; M is hydrogen, F, Cl or Br; R is F, Cl, Br, CF₃ or OCF₃; W is NO₂ and X and Y are each independently Cl, Br or CF₃.

Illustrative of some of the insecticidal, acaricidal and nematocidal arylpyrroles of the present invention are:

- 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile;
- 10 4,5-dichloro-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile;
- 4-bromo-5-chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile;
- 5-bromo-4-chloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile;
- 4,5-dichloro-2-(o-chlorophenyl)pyrrole-3-carbonitrile;
- 2-(p-bromophenyl)-4,5-dichloropyrrole-3-carbonitrile;
- 15 4,5-dichloro-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
- 4,5-dibromo-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
- 4,5-dibromo-2-(o-chlorophenyl)pyrrole-3-carbonitrile;
- 4,5-dibromo-2-(p-chlorophenyl)pyrrole-3-carbonitrile;
- 4,5-dichloro-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile;
- 20 4,5-dibromo-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile;
- 2,3-dibromo-4-nitro-5-phenylpyrrole;
- 2-(p-bromophenyl)-4,5-dichloro-3-nitropyrrole;
- 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl)-pyrrole;
- 4,5-dichloro-2-(m-chlorophenyl)pyrrole-3-carbonitrile;
- 25 4,5-dichloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile;
- 4,5-dichloro-2-phenylpyrrole-3-carbonitrile;
- 2,3-dichloro-5-(p-chlorophenyl)-4-nitropyrrole;
- 2-bromo-3-chloro-5-(p-chlorophenyl)-4-nitropyrrole;
- 2,3-dibromo-5-(p-chlorophenyl)-4-nitropyrrole;
- 30 2,3-dichloro-4-nitro-5-phenylpyrrole;
- 3-bromo-2-chloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl)-pyrrole;
- 5-Chloro-2-(3,4-dichlorophenyl)-1-(methoxymethyl)-4-(trifluoromethyl)pyrrole-3-carbonitrile;
- 5-Bromo-2-(m-fluorophenyl)-3-nitro-4-(trifluoromethyl) pyrrole;
- 2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
- 35 3-Bromo-5-(m-fluorophenyl)-4-nitro-2-(trifluoromethyl) pyrrole;
- 4-Bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
- 4-Chloro-2-(3,5-dichloro-4-methylphenyl)-3-nitro-5-(trifluoromethyl)pyrrole;
- 2-(2-Bromo-4-chlorophenyl)-1-(2-propynyl)-4,5-bis-(trifluoromethyl)pyrrole-3-carbonitrile;
- 2-(2,5-Difluorophenyl)-3-nitro-4,5-bis-(trifluoromethyl)pyrrole;
- 40 5-[p-(Trifluoromethoxy)phenyl]pyrrole-2,4-dicarbonitrile;
- 5-(p-Dimethylaminophenyl)-4-nitropyrrole-2-carbonitrile;
- 3-Bromo-5-(p-chlorophenyl)pyrrole-2,4-dicarbonitrile;
- 4-Bromo-2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile;
- 5-(p-Methylthiophenyl)-3-(trifluoromethyl)pyrrole-2,4-dicarbonitrile;
- 45 1-Allyl-4-nitro-5-(α,α,α -trifluoro-p-tolyl)-3-(trifluoromethyl)pyrrole-2-carbonitrile;
- 4-Chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile;
- 2-(m-Methanesulfonylphenyl)-4-(trifluoromethyl)pyrrole -3-carbonitrile;
- 2-(3-chloro-4-methylphenyl)-1-methyl-3-nitro-4-(trifluoromethyl)pyrrole;
- 2-Phenylpyrrole-3,4-dicarbonitrile;
- 50 5-(p-Ethanesulfinylphenyl)-4-nitropyrrole-3-carbonitrile;
- 2-Bromo-5-phenylpyrrole-3,4-dicarbonitrile;
- 2-Chloro-5-(3,5-dichlorophenyl)-4-nitropyrrole-3-carbonitrile;
- 1-Benzyl-4-nitro-5-(p-chlorophenyl)-2-(trifluoromethyl) pyrrole-3-carbonitrile;
- 2-Chloro-5-(m-bromophenyl)pyrrole-3-carbonitrile;
- 55 2-Bromo-1-(p-chlorophenoxy)methyl-5-(p-chlorophenyl) -3-nitropyrrole;
- 2,4-Dibromo-5-phenylpyrrole-3-carbonitrile;
- 5-(p-Bromophenyl)-2,4-dichloro-3-nitropyrrole;
- 2-Bromo-5-(3-bromo-4-methylphenyl)-1-(n-propyloxy) methyl-4-(trifluoromethyl)pyrrole-3-carbonitrile;


- 2-Bromo-5-(p-chlorophenyl)-3-nitro-4-(trifluoromethyl) pyrrole;
 5-[m-(Difluoromethoxy)phenyl]-2-(trifluoromethyl) pyrrole-3-carbonitrile;
 5-(2,3-Dichlorophenyl)-1-methoxymethyl-3-nitro-2-(trifluoromethyl)pyrrole;
 4-Chloro-5-(β -naphthyl)-2-(trifluoromethyl)pyrrole-3-carbonitrile;
 5 3-Bromo-2-(3,4-dichlorophenyl)-4-nitro-5-(trifluoromethyl)pyrrole;
 5-(2-Bromo-5-ethylphenyl)-2,4-bis-(trifluoromethyl) pyrrole-3-carbonitrile;
 1-Ethyl-2-(p-fluorophenyl)-4-nitro-3,5-bis-(trifluoromethyl)pyrrole;
 1-[(2,6-Dichlorophenoxy)methyl]-5-(m-chlorophenyl) pyrrole-2,3-dicarbonitrile;
 3-Nitro-5(α,α,α -trifluoro-p-tolyl)pyrrole-2-carbonitrile;
 10 4-Chloro-5-(4-chloro-2-methylphenyl)pyrrole-2,3-dicarbonitrile;
 4-Bromo-5-(3,4-dibromophenyl)-2-nitropyrrole-3-carbonitrile;
 1-[(1-Methoxy)ethyl]-5-(p-chlorophenyl)-4-(trifluoromethyl)pyrrole-2,3-dicarbonitrile;
 5-(p-Isopropylphenyl)-2-nitro-4-(trifluoromethyl) pyrrole-3-carbonitrile;
 4-Chloro-5-(3,4-difluoromethylenedioxyphenyl)pyrrole-3-carbonitrile;
 15 3-Bromo-2-(3-chloro-4-cyanophenyl)-4-nitropyrrole;
 1-[(3,4-dichlorobenzoyloxy)methyl]-2-(m-bromophenyl) pyrrole-4-carbonitrile;
 2-(3,5-Dichloro-4-methylphenyl)-4-nitro-3-trifluoromethylpyrrole;
 2-Phenylpyrrole-3,4-dicarbonitrile;
 2-(2-Bromo-4-chlorophenyl)-4-nitropyrrole-3-carbonitrile;
 20 2-Bromo-5-phenylpyrrole-3,4-dicarbonitrile;
 5-Chloro-2-(3,4-dibromophenyl)-1-methyl-4-nitropyrrole-3-carbonitrile;
 2-(p-Chlorophenyl)-5-(trifluoromethyl)pyrrole-3,4-dicarbonitrile;
 2-(o-Bromophenyl)-4-nitro-5-(trifluoromethyl)pyrrole-3-carbonitrile;
 3-Bromo-5-(3-chloro-4-methoxy)pyrrole-2-carbonitrile;
 25 3-Bromo-5-(m-bromophenyl)-2-nitropyrrole;
 3,4-Dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile;
 2-(3-Chloro-4-cyanophenyl)-5-nitro-3,4-dichloropyrrole;
 3-Chloro-1-(p-methoxybenzyl)-5-(3,4-difluorophenyl)-4-(trifluoro-methyl)pyrrole-2-carbonitrile;
 3-Bromo-5-(3,5-dibromo-p-tolyl)-2-nitro-4-(trifluoromethyl)pyrrole;
 30 1-(2,3,3-Trichloroallyl)-5-(p-chlorophenyl)-3-(trifluoromethyl)pyrrole-2-carbonitrile;
 2-(p-Iodophenyl)-5-nitro-4-(trifluoromethyl)pyrrole;
 4-Chloro-5-(m-isopropylphenyl)-3-(trifluoromethyl) pyrrole-2-carbonitrile;
 3-Bromo-1-methyl-2-(3-fluoro-4-methylphenyl)-2-nitro-3-(trifluoromethyl)pyrrole;
 5-(p-Bromophenyl)-1-isopropyl-3,4-bis-(trifluoromethyl) pyrrole-2-carbonitrile;
 35 2-(3,4-Dichloro-4-methylthio)-5-nitro-3,4-bis-(trifluoromethyl)pyrrole;
 5-(m-Difluoromethoxyphenyl)pyrrole-2,3-dicarbonitrile;
 5-(3-Bromo-4-cyanophenyl)-2-nitropyrrole-3-carbonitrile;
 4-Chloro-1-methoxymethyl-5-(p-bromophenyl)pyrrole-2,3-dicarbonitrile;
 4-Bromo-5-(2,6-dichloro-4-methylthio)-2-nitropyrrole-3-carbonitrile;
 40 1-[(p-Bromophenoxy)methyl]-5-(m-trifluoromethyl)-4-(trifluoromethyl)pyrrole-2,3-dicarbonitrile;
 5-(α -Naphthyl)-2-nitro-4-(trifluoromethyl)pyrrole-3-carbonitrile;
 4-Bromo-5-(3-bromo-4-trifluoromethylphenyl)pyrrole-2-carbonitrile;
 3-Chloro-2-(2,3-dichlorophenyl)-5-nitropyrrole;
 5-(m-Cyanophenyl)-3-(trifluoromethyl)pyrrole-2-carbonitrile;
 45 2-(3-Bromo-4-isopropoxy)-5-nitro-3-(trifluoromethyl) pyrrole;
 5-(p-Chlorophenyl)pyrrole-2,4-dicarbonitrile;
 2-(3,4-Dichlorophenyl)-5-nitropyrrole-3-carbonitrile;
 3-Bromo-5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonitrile;
 4-Bromo-2-(3,4-dichlorophenyl)-5-nitropyrrole-3-carbonitrile;
 50 5-(3,4-Dibromophenyl)-3-(trifluoromethyl)pyrrole-2,4-dicarbonitrile;
 2-(m-Chlorophenyl)-5-nitro-4-(trifluoromethyl)pyrrole-3-carbonitrile;
 5-Bromo-3-(3,5-dichloro-4-difluoromethoxyphenyl) pyrrole-2-carbonitrile;
 2-Bromo-4-(2,5-dibromophenyl)-5-nitropyrrole;
 2,3-Dibromo-4-(p-chlorophenyl)pyrrole-5-carbonitrile;
 55 2,3-Dichloro-4-(3,5-difluorophenyl)-5-nitropyrrole;
 5-Bromo-3-(p-chlorophenyl)-1-hydroxyethyl-4-(trifluoromethyl)pyrrole-2-carbonitrile;
 2-Chloro-5-nitro-3-(trifluoromethyl)-4-(m-trifluoromethylphenyl)pyrrole;
 3-(3-Bromo-4-chlorophenyl)-5-(trifluoromethyl)pyrrole-2-carbonitrile;

- 3-(3-Chloro-4-fluorophenyl)-2-nitro-5-(trifluoromethyl) pyrrole;
 4-Bromo-3-(p-chlorophenyl)-1-methylthiomethyl-5-(trifluoromethyl)pyrrole-2-carbonitrile;
 3-(4-Bromo-3-cyanophenyl)-4-chloro-2-nitro-5-(trifluoromethyl)pyrrole;
 4-(p-Chlorophenyl)-2,3-bis-(trifluoromethyl)pyrrole-2-carbonitrile;
 5 3-(2,3-Dichlorophenyl)-2-nitro-4,5-bis-(trifluoromethyl)pyrrole;
 3-(3,4-Dichlorophenyl)pyrrole-2,5-dicarbonitrile;
 4-(2-Bromo-4-methylphenyl)-5-nitropyrrole-2-carbonitrile;
 3-Bromo-4-(3,5-dichloro-4-methylthiophenyl)pyrrole-2,5-dicarbonitrile;
 4-(m-Bromophenyl)-3-chloro-5-nitropyrrole-2-carbonitrile;
 10 3-(p-Acetamidophenyl)-4-(trifluoromethyl)pyrrole-2,5-dicarbonitrile;
 4-(m-Bromophenyl)-5-nitro-3-(trifluoromethyl)pyrrole-2-carbonitrile;
 4-Chloro-3-(3,4-dichlorophenyl)-1-(1-propenyl)pyrrole-2-carbonitrile;
 3-Bromo-4-(p-dimethylaminophenyl)-5-nitropyrrole;
 1-(3,4-Dichlorobenzyl)-3-(p-chlorophenyl)-4-(trifluoromethyl)pyrrole-2-carbonitrile;
 15 2-Nitro-3-(p-tetrafluoroethoxyphenyl)-4-(trifluoromethyl)pyrrole;
 3-(3-Bromo-4-*i*-propylphenyl)pyrrole-2,4-dicarbonitrile;
 4-(p-Ethylsulfonylphenyl)-5-nitropyrrole-3-carbonitrile;
 5-Bromo-1-(2-methoxyethyl)-4-(2,4,6-trichlorophenyl) pyrrole-2,4-dicarbonitrile;
 2-Chloro-4-(2,3-dichlorophenyl)-5-nitropyrrole-3-carbonitrile;
 20 3-(p-Fluorophenyl)-5-(trifluoromethyl)pyrrole-2,4-dicarbonitrile;
 4-(p-Iodophenyl)-5-nitro-2-(trifluoromethyl)pyrrole-3-carbonitrile;
 5-Chloro-4-[p-(N-methylacetamido)phenyl]pyrrole-2-carbonitrile;
 5-Bromo-4-(o-bromophenyl)-1-propargylpyrrole-2-carbonitrile;
 2-Bromo-3-(o-bromophenyl)-5-nitropyrrole;
 25 4-(p-Chlorophenyl)-3,5-dichloro-1-(2,3,3-trichloroallyl) pyrrole-2-carbonitrile;
 3-Bromo-5-chloro-4-(p-chlorophenyl)-2-nitropyrrole;
 5-Bromo-4-[p-(2,2-dichloro-1,1-difluoroethoxy)phenyl] -3-(trifluoromethyl)pyrrole-2-carbonitrile;
 2-Chloro-3-(2-bromo-4-ethylthiophenyl)-5-nitro-4-(trifluoromethyl)pyrrole;
 3-(3-Bromo-4-acetylphenyl)-5-(trifluoromethyl)pyrrole-2-carbonitrile;
 30 1-Cyano-3-(3,4-dibromophenyl)-5-nitro-2-(trifluoromethyl)pyrrole;
 3-Bromo-1-methoxymethyl-4-(m-trifluoromethyl)-5-(trifluoromethyl)pyrrole-2-carbonitrile;
 3-(p-Chlorophenyl)-4-iodo-5-nitro-2-(trifluoromethyl) pyrrole;
 4-(p-Bromophenyl)-1-[(1-ethoxy)ethyl]-3,5-di-(trifluoromethyl)pyrrole-2-carbonitrile;
 3-(2-Bromo-4-methoxyphenyl)-5-nitro-2,4-di-(trifluoromethyl)pyrrole;
 35 3-(p-Chlorodifluoromethoxyphenyl)pyrrole-2,5-dicarbonitrile;
 2-(p-Isobutyrylamino)phenyl)-5-nitropyrrole-2-carbonitrile;
 3-Bromo-4-(3,4-dimethoxyphenyl)pyrrole-2,5-dicarbonitrile;
 4-Chloro-3-(p-chlorophenyl)-1-isopropoxyloxycarbonylmethyl)-5-nitropyrrole-2-carbonitrile;
 3-(o-Bromophenyl)-4-(trifluoromethyl)pyrrole-2,5-dicarbonitrile;
 40 1-(2-Chloroethyl)-3-(3,4-dichlorophenyl)-4-(trifluoromethyl)pyrrole-2-carbonitrile;
 4-(4-Bromo-3-trifluoromethoxyphenyl)-3-chloropyrrole-2-carbonitrile;
 3-Bromo-4-(2,4-dichlorophenyl)-1-isopropyl-2-nitropyrrole;
 4-(3-Methoxy-4-cyanophenyl)-3-(trifluoromethyl)pyrrole -2-carbonitrile;
 1-(3,4-Dichlorobenzyl)-4-(2-methyl-4-iodophenyl)-2-nitro-3-trifluoromethylpyrrole;
 45 1-Methyl-4-[3,5-di(trifluoromethyl)phenyl]pyrrole-2,3-dicarbonitrile;
 4-(3,4-Dichlorophenyl)-2-nitropyrrole-3-carbonitrile;
 4-(m-Bromophenyl)-1-carbomethoxymethyl-5-chloropyrrole-2,3-dicarbonitrile;
 5-Bromo-4-(2,6-dichloro-4-methanesulfinylphenyl)-2-nitropyrrole-3-carbonitrile;
 4-(p-Chlorophenyl)-1-(2,2,2-trifluoroethyl)-5-(trifluoromethyl)pyrrole-2,3-dicarbonitrile;
 50 4-(3,5-Dichlorophenyl)-2-nitro-5-(trifluoromethyl) pyrrole-3-carbonitrile;
 2-Chloro-4-(3-chloro-4-N-methylacetamidophenyl) pyrrole-3-carbonitrile;
 2-Bromo-4-(3-bromo-4-n-propylphenyl)-3-nitropyrrole;
 2,5-Dichloro-4-(3,5-dichloro-4-methylthiophenyl) pyrrole-3-carbonitrile;
 2,5-Dibromo-1-(2,4-dibromophenoxy)methyl)-3-(p-chlorophenyl)-4-nitropyrrole;
 55 4-(3-Bromo-4-cyanophenyl)-2-chloro-5-(trifluoromethyl) pyrrole-3-carbonitrile;
 2-Bromo-1-methyl-3-nitro-4-(α,α,α -trifluoro-p-tolyl) pyrrole;
 4-(p-chlorophenyl)-1-(n-butyloxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
 4-(3,4-Methylenedioxyphenyl)-3-nitro-2-(trifluoromethyl)pyrrole;

- 5-Chloro-4-(3-chloro-4-trifluoromethoxyphenyl)-2-(trifluoromethyl)pyrrole-3-carbonitrile;
 2-Bromo-3-(3,4-dichlorophenyl)-1-ethylthiomethyl-4-nitro-5-(trifluoromethyl)pyrrole;
 4-[p-(tetrafluoroethoxy)phenyl]-2,5-di-(trifluoro methyl)pyrrole-3-carbonitrile;
 3-(3-Bromo-4-acetoxyphenyl)-1-(3,4-dichlorophenoxymethyl)-4-nitro-2,5-di-(trifluoromethyl) pyrrole;
 5 4-[p-Bromophenyl]-1-[(2-methoxy)ethyl]pyrrole-2,3-dicarbonitrile;
 4-(m-Isopropionamidophenyl)-3-nitropyrrole-2-carbonitrile
 5-Bromo-4-(2-chloro-4-methylthio)pyrrole-2,3-dicarbonitrile;
 5-Chloro-4-(p-chlorophenyl)-1-hydroxyethyl-3-nitropyrrole-2-carbonitrile;
 4-(3,5-Dibromo-4-cyanophenyl)-5-(trifluoromethyl) pyrrole-2,3-dicarbonitrile;
 10 4-(4-Chloro-2-methylphenyl)-1-isopropylthiomethyl-3-nitro-5-(trifluoromethyl)pyrrole-2-carbonitrile;
 5-Bromo-4-(3,4-dichlorophenyl)-1-(difluoromethyl) pyrrole-3-carbonitrile;
 2-Chloro-3-(m-difluoromethoxyphenyl)-4-nitropyrrole;
 1-(2,4-Dibromophenoxymethyl)-4-(m-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
 3-(3-Bromo-4-ethoxy)-4-nitro-2-(trifluoromethyl) pyrrole;
 15 3-(2,4,6-Trichlorophenyl)pyrrole-2,4-dicarbonitrile;
 3-(4-Bromo-3-chlorophenyl)-1-(difluoromethyl)-4-nitropyrrole-2-carbonitrile;
 5-Bromo-3-(p-chlorophenyl)-1-(isobutyloxymethyl) pyrrole-3-carbonitrile;
 3-(4-Bromo-3-methylphenyl)-5-chloro-4-nitropyrrole-2-carbonitrile;
 3-(2-Naphthyl)-5-(trifluoromethyl)pyrrole-2,4-dicarbonitrile;
 20 3-(3-Cyano-4-methylphenyl)-1-methyl-4-nitro-5-(trifluoromethyl)pyrrole-2-carbonitrile;
 2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole
 2-(3,5-dibromo-4-methoxyphenyl)-4,5-dichloropyrrole-3-carbonitrile;
 2,3-dichloro-4-nitro-5-(2,4,6-trifluorophenyl)-4-nitropyrrole;
 4,5-dibromo-2-(2,3,6-trifluorophenyl)-3-carbonitrile
 25 4,5-dichloro-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)-pyrrole-3-carbonitrile;
 4,5-dibromo-1-methyl-2-(α,α,α -trifluoro-p-tolyl) pyrrole-3-carbonitrile;
 4,5-dichloro-2-(3,4-dichlorophenyl)-1-ethylpyrrole-3-carbonitrile;
 2,3-dichloro-4-nitro-5-[p-(trifluoromethoxy)phenyl]pyrrole;
 4,5-dichloro-2-[m-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile;
 30 4,5-dichloro-2-(3,4-dichlorophenyl)-1-methylpyrrole-3-carbonitrile;
 2,3-dichloro-5-(p-chlorophenyl)-1-methyl-4-nitro pyrrole;
 and
 4-bromo-5-chloro-2-(p-chlorophenyl)-1-methylpyrrole-3-carbonitrile.
 5-chloro-2-(3,4-dichlorophenyl)-4-fluoropyrrole-3-carbonitrile
 35 2-bromo-5-(p-chlorophenyl)-1-(ethoxymethyl)-4-fluoropyrrole-3-carbonitrile
 3-bromo-5-(p-chlorophenyl)-2-fluoro-4-nitropyrrole
 The α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -cyanostyrene and α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene compounds of this invention are depicted by the following structural formula:


40

45



- 50 wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:
 -OCH₂O-, -OCF₂- or

55



5

Z is S(O)*n* or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and *n* is an Integer of 0, 1 or 2.

A preferred group of β-(substituted)styrene compounds of the present invention have the above-illustrated structure wherein W is CN; L is H, Cl or Br; M is H, F, Cl, Br or OCH₃; R is H, F, Cl, Br, CF₃, NO₂, OCF₃ or OCH₃; or when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represents the structure:

15



Another preferred group of β-(substituted)styrene compounds of this invention have the above-illustrated structure wherein W is NO₂; L is H, Cl or Br; M is H, F, Cl, Br or OCH₃; R is H, F, Cl, Br, CF₃, NO₂, OCF₃ or OCH₃; or when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represents the structure:

25




30

While the compounds of the present invention are referred to above as β-cyanostyrenes and β-nitrostyrenes, they may also be named as dialkyl acetals.

Some of the preferred dialkyl acetal compounds of this invention are (E) and (Z) (1)p-chloro-β-[(formylmethyl)amino]cinnamonnitrile diethyl acetal; (2)β-[(formylmethyl)amino]-3,4-dimethoxycinnamonnitrile diethyl acetal; (3)(Z)-methyl p-(2-cyano-1-[(formylmethyl)amino]vinyl)benzoate diethyl acetal; (4)(Z)-β-[(formylmethyl)amino]-1-naphthaleneacrylonitrile diethyl acetal; (5)(Z)-β-[(formylmethyl)amino]-p-methylcinnamonnitrile diethyl acetal; (6)N-(formylmethyl)-p-methyl-α-(nitromethylene)benzylamine diethyl acetal; (7)N-(formylmethyl)-3,4-dimethoxy-α-(nitromethylene)benzylamine diethyl acetal; (8)N-(formylmethyl)-α-(nitromethylene)-2-naphthalenemethylamine diethyl acetal; (9)methyl p-(α-[(formylmethyl)amino]-β-nitrovinyl)benzoate p-(diethyl acetal); (10)N-(formylmethyl)-3,4-dimethoxy-α-(nitromethylene)benzylamine dimethyl acetal; (11)(E) and (Z) p-chloro-β-[(formylmethyl)amino]cinnamonnitrile dimethyl acetal; (12)β-[(formylmethyl)amino]-3,4-dimethoxycinnamonnitrile dimethyl acetal; (13)3,4-dichloro-β-[(formylmethyl)amino]cinnamonnitrile diethyl acetal; and (14)p-trifluoromethyl-β-[(formylmethyl)amino]cinnamonnitrile diethyl acetal.


The β-cyanostyrenes, also referred to as cinnamonnitrile dialkyl acetals, can be prepared by the reaction of a substituted or unsubstituted benzoyl acetonitrile with a 2,2-di(C₁-C₄ alkoxy)ethylamine in the presence of an aromatic solvent to form the α-(2,2-di(C₁-C₄ alkoxy)ethylamino)-β-cyano-(substituted)styrene which then may be converted to a 2-(substituted-phenyl)pyrrole-3-carbonitrile by reaction of said β-3-cyano-(substituted)styrene compound with trifluoroacetic acid. Chlorination of the thus prepared cyanophenyl pyrrole with sodium hypochlorite or sulfuryl chloride in an inert solvent yields the insecticidal, acaricidal, and nematocidal 4,5-dichloro-2-(substituted-phenyl)pyrrole-3-carbonitrile. The conversion to the pyrrole intermediate may also be achieved by substituting concentrated HCl at a temperature between about 20 and 40 °C. The reactions may be graphically illustrated as follows:

55




wherein A is hydrogen, C₁-C₄ alkyl optionally substituted with one C₁-C₄ alkoxy, one C₁-C₄ alkylthio, from one to three halogen groups, or phenyl optionally substituted with one or two C₁-C₄ alkyl, C₁-C₄ alkoxy, or halogen groups; C₃-C₄ alkenyl optionally substituted with from to three halogen groups; or C₃-C₄ alkynyl; X is Cl or Br; R₆ is C₁-C₄ alkyl and L, R and M are as described above.

Certain novel arylpyrrole compounds of formula I, wherein A is hydrogen; W is CN and X, Y, L, M and R are as described above, can be prepared by reacting N-formyl-DL-phenyl-glycine or a substituted N-formylphenylglycine represented by the structure formula VIII:



wherein L is H, F, Cl or Br; R and M are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached, M and R may form a ring in which MR represents the structure:


-OCH₂O-, -OCF₂O- or




;

Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCI or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl or R₅CO; R₅ is H or C₁-C₃ alkyl and n is an integer of 0, 1 or 2; with at least an equivalent amount of a 2-chloroacrylonitrile and two to three equivalents of acetic anhydride. The reaction is conducted at an elevated temperature, preferably about 70° to 100°C.


The reaction can be illustrated as follows:



Conversion of the thus prepared 2-phenylpyrrole-3-carbonitrile or 2-(substituted phenyl)pyrrole-3-carbonitrile to the corresponding formula II, 4-halo, 5-halo or 4,5-dihalo-2-(substituted phenyl)pyrrole-3-carbonitrile, is readily achieved by reaction of the above said 2-phenylpyrrole-3-carbonitrile or 2-(substituted phenyl)pyrrole-3-carbonitrile with at least about 1 or 2 equivalents of a sulfuryl halide, bromine or chlorine, in the presence of a solvent such as dioxane, THF, acetic acid or a chlorinated hydrocarbon solvent. For preparation of a monohalo pyrrole-3-carbonitrile use of about 1 equivalent of the halogenating agent is required; whereas, preparation of a dihalo pyrrole-3-carbonitrile requires 2 to 3 equivalents of said halogenating agent. When sulfuryl chloride or sulfuryl bromide is used the reaction is generally conducted at a temperature below about 40°C and preferably between about 0° and 30°C, but when elemental bromine is employed, the reaction is usually conducted at about 30-40°C. Other effective halogenating agents that may be employed in these reactions include sodium hypochlorite, *t*-butylhypochlorite, N-bromosuccinimide, N-iodosuccinimide and the like. The reaction may be illustrated as follows:



The formula II carbonitrile compounds of the present invention may also be prepared from the reaction of a substituted or unsubstituted benzoyl acetonitrile with a 2,2-di(C₁-C₄ alkoxy)ethylamine in the presence of an aromatic solvent to form the α -(2,2-di(C₁-C₄ alkoxy)ethylamino)- β -cyano-(substituted)styrene which is then converted to the 2-(substituted-phenyl)pyrrole-3-carbonitrile of formula II by reaction of said β -3-cyano-(substituted)styrene compound with trifluoroacetic acid or with concentrated HCl at a temperature between about 20° and 40°C. The reactions may be graphically illustrated as follows:




50 wherein R₆ is C₁-C₄ alkyl and L, R and M are as described above.

Also in accordance with the present invention formula II 3-nitro-2-phenylpyrrole and 3-nitro-2-(substituted)phenylpyrrole compounds can be prepared by reaction of an α -nitroacetophenone or a substituted α -nitroacetophenone with a 2,2-di(C₁-C₄-alkoxy)ethylamine. The reaction is generally conducted in the presence of an inert organic solvent preferably an aromatic solvent, at an elevated temperature to give an α -(2,2-di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrene or a substituted α -(2,2-di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrene that is converted to the formula II 3-nitro-2-phenylpyrrole or 3-nitro-2-(substituted)phenylpyrrole by treatment with a mineral acid such as hydrochloric or hydrobromic acid. Reaction of the thus prepared nitrophenylpyrrole with sodium hypochlorite in the presence of an inert organic solvent at a reduced


temperature yields the formula II 2,3-dichloro-4-nitro-5-phenyl or 5-(substituted)phenylpyrrole.

The above reactions may be graphically illustrated as follows:




35 In addition to the several methods described in the literature for preparing substituted and unsubstituted benzoyl acetonitriles, surprisingly we have found that these compounds may also be prepared by reacting an appropriately substituted benzoyl halide with an alkali metal hydride and an alkyl cyanoacetate, such as *t*-butyl cyanoacetate, to yield the corresponding *t*-butyl(benzoyl or substituted benzoyl)cyanoacetate. These

40 reactions may be graphically illustrated as follows:




The thus formed cyanoacetate ester can then be converted to a substituted or unsubstituted benzoyl acetonitrile by heating the compound in toluene containing *p*-toluene sulfonic acid. The reaction may be graphically illustrated as follows:



10 Examples of the *t*-butyl(benzyl and substituted benzyl) acetonitriles used in the above reactions are shown in Tables below.

t-Butyl(benzyl and Substituted benzyl)cyanoacetates




30

L	M	R	mp °C
H	3-Cl	4-Cl	91-94
H	H	4-OCF ₃	81-84
H	H	4-Br	113-115
35	H	4-CF ₃	146-147
H	H	4-F	98-100
H	H	4-CN	127-128
H	H	4-CF ₃ CH ₂ O	136-139
H	H	4-CH ₃ SO ₂	127-129
40	H	3-F	91-94
H	H	4-CH ₃ S	117-119.5
H	H	4-CHF ₂ CF ₂ O	92-94
3-Cl	5-Cl	4-CH ₃ O	---

45


50

55

Benzoyl Acetonitriles


L	M	R	mp °C
H	H	4-Cl	128.5-129.5
H	3-Cl	4-Cl	105-107
H	H	2-C	153-55
H	H	4-OCF ₃	79-81
H	H	4-CF ₃	44-45
H	2-Cl	4-Cl	66-67
H	H	3-Cl	80-83
H	H	4-CN	126-128
H	H	4-F	78-80
H	H	4-SO ₂ CH ₃	129-132
H	3-F	4-F	74-75
H	H	3-CF ₃	58-60
H	H	4-CH ₃	103.5-106
H	H	4-NO ₂	119-124
3-Cl	5-Cl	4-OCH ₃	---

Preparation of N-substituted formula I arylpyrroles can be achieved by reaction of the appropriately substituted formula I arylpyrrole, wherein A is hydrogen and L, M, R, W, X and Y are as described above, with an appropriate alkylating agent and a suitable base. For example, a brominated hydroxy-C₁-C₄-alkyl and potassium t-butoxide. This reaction provides an arylpyrrole having the same substituents as the starting material, but in addition is substituted on the nitrogen with hydroxy-C₁-C₄ alkyl. In a similar reaction cyanogen bromide is substituted for the brominated hydroxy C₁-C₄ alkyl and yields the formula I arylpyrrole with a carbonitrile substituent on the nitrogen. The reactions may be illustrated as follows:



wherein L, M, R, W, X and Y are as described for formula I above and A is 1) C₁-C₄ alcohol or 2) CN.

20 Preparation of 2-phenylpyrrole 3,4-dicarbonitrile, 2-bromo-5-phenylpyrrole-3,4-dicarbonitrile and substituted phenyl derivatives thereof can be obtained by reaction of fumaronitrile with bromine in the presence of a chlorinated hydrocarbon such as chloroform at an elevated temperature to yield bromofumaronitrile. The thus formed bromofumaronitrile is then reacted with N-(trimethylsilyl)methyl-5-methyl-benzene-thioimide or a substituted derivative thereof, in the presence of hexamethylphosphoramide at an elevated temperature to yield the 2-phenylpyrrole-3,4-dicarbonitrile. Bromination of the thus prepared 3,4-dicarbonitrile yields the 2-bromo-5-phenylpyrrole-3,4-dicarbonitrile or the substituted phenyl derivative if the substituted N-(trimethylsilyl)methyl-5-methyl-benzene-thioimide is used in the previous reaction. The reaction may be graphically illustrated as follows:



The examples provided by way of illustration below utilize the schemes illustrated above and provide a means for preparing other compounds of the invention which are not specifically described herein.

55 The arylpyrroles of the present invention are effective for controlling insects, acarina and nematodes. These compounds are also effective for protecting growing or harvested crops from attack by the above-said pests.

In practice generally about 10 ppm to about 10,000 ppm and preferably 100 to about 5000 ppm, of the formula I arylpyrrole, which encompasses all of the arylpyrrole isomers of formulas II, III, IV, V, VI and VII,

dispersed in water or other inexpensive liquid carrier is effective when applied to the plants, the crops or the soil in which said crops are growing to protect said crops from attack by insects, acarina and/or nematodes. These compounds are also useful for protecting turf grass from attack by pests such as grubs, chinch bugs and the like.

5 The formula I arylpyrroles of this invention are also effective for controlling insects, nematodes and acarina, when applied to the foliage of plants and/or to the soil or water in which said plants are growing in sufficient amount to provide a rate of from about 0.125 kg/ha to about 4.0 kg/ha of active ingredient. Obviously higher rates of application of the formula I arylpyrroles may be used to protect crops from attack by insects, nematodes and acarina, however, higher rates of application are generally unnecessary and
10 wasteful.

While the arylpyrroles of this invention are effective for controlling insects, nematodes and acarina when employed alone, they may be used in combination with other biological chemicals, including other insecticides, nematocides and acaricides. For example, the arylpyrroles of this invention may be used effectively in conjunction or combination with phosphates, carbamates, pyrethroids, formamides, chlorinat-
15 ed hydrocarbons, halobenzoyleureas and the like.

The 2-aryl-3-cyano-4,5-dihaloarypyrroles prepared from the β -cyano-styrene compounds of the present invention are effective for controlling insects, acarina and nematodes. These compounds are also effective for protecting growing or harvested crops from attack by the above-said pests.

In practice generally about 10 ppm to 10,000 ppm and preferably 100 to 5000 ppm, of the halogenated
20 arylpyrrole dispersed in water or other inexpensive liquid carrier is effective when applied to the plants, the crops or the soil in which said crops are growing to protect said crops from attack by insects, acarina and/or nematodes.

The above-said halogenated arylpyrroles are also effective for controlling insects, nematodes and acarina, when applied to the foliage of plants and/or to the soil or water in which said plants are growing. These halogenated arylpyrrole compounds are usually applied in sufficient amount to provide a rate of from
25 about 0.125 kg/ha to about 4.0 kg/ha of active ingredient. Obviously higher rates of application of said halogenated arylpyrroles may be used to protect crops from attack by insects, nematodes and acarina, however, higher rates of application are generally unnecessary and wasteful.

Advantageously, the above-said arylpyrroles may be formulated into dry compacted granules, flowable
30 compositions, granular formulations, wettable powders, emulsifiable concentrates, dusts, dust concentrates, microemulsions and the like, all of which lend themselves to soil, water and/or foliage application and provide the requisite plant protection. Such formulations include the compounds of the invention admixed with inert, pharmacologically- acceptable solid or liquid diluents.

For example, wettable powders, dusts and dust concentrate formulations of the invention can be
35 prepared by grinding together about 3% to 20%, by weight, of the formula I arylpyrrole compound, with about 3% to 20% by weight of a solid anionic surfactant. One suitable anionic surfactant is a dioctyl ester of sodium sulfosuccinic acid, specifically Aerosol OT® surfactant marketed by the American Cyanamid Company. About 60% to 94%, by weight, of an inert solid diluent, such as montmorillonite, attapulgite, chalk, talc, kaolin, diatomaceous earth, limestone, silicates or the like also is used in such formulations.

40 Compacted granules especially useful for soil or water application can be prepared by grinding together in about equal parts, usually about 3 to 20 parts, of the arylpyrrole and a solid surfactant, with about 60 to 94 parts of gypsum. Thereafter, the mixture is compacted into small granular particles, about 24/48 mesh or larger.

Other suitable solid surfactants useful in the present formulations include not only the anionic dioctyl
45 ester of sodium sulfosuccinic acid but also nonionic block copolymers of ethylene oxide and propylene oxide. Such block copolymers are marketed by BASF Wyandotte Corporation as Pluronic 10R8®, 17R8®, 25R8®, F38®, F68®, F77® or F87®, and are especially effective for the preparation of compacted granules.

In addition to the powders and concentrate formulations described hereinabove, wettable powders and flowables may be used because they may be dispersed in water. Preferably, such flowables will be applied
50 at the locus with the aqueous compositions being sprayed on the foliage of plants to be protected. These sprays also may be applied to the breeding ground, food supply or habitat of the insects and acarina sought to be controlled.


Where solid formulations of the arylpyrroles are to be used in combination treatments with other
55 pesticidal agents, the formulations can be applied as an admixture of the components or may be applied sequentially.

Similarly, liquid formulations of the arylpyrrole in combination with other pesticidal agents may be tank mixed or may be applied separately, sequentially, as liquid sprays. Liquid spray formulations of the compounds of the invention should contain about 0.001% to 0.1% by weight of the active arylpyrrole.

The following examples are presented as illustrations of the present invention.

EXAMPLE 1

2-Phenylpyrrole-3-carbonitrile




The following procedure is similar to the method given in JOC, 43, 4273-6 (1978). A magnetically stirred mixture of 30.00g of N-formyl-phenylglycine is heated at 90 °C for 1 & 1/2 hours. The clear yellow reaction solution is concentrated in vacuo to give 42.5g of an oily brownish orange semi-solid. Material partially purified by chromatography on silica gel is shown by the proton NMR spectrum to be a mixture of 73% 2-phenylpyrrole-3-carbonitrile and 27% 2-phenyl-3-cyano-5-methylpyrrole. Recrystallization once from chloroform and twice from 1,2-dichloroethane gives 1.69g of an off-white solid which proton NMR shows it to be 96% 2-phenylpyrrole-3-carbonitrile, mp 148-152 °C.

Microanalysis (MW 168.19):

Calcd.:	C, 78.55%;	H, 4.79%;	N, 16.66%
Found:	C, 78.52%;	H, 4.73%;	N, 16.54%

EXAMPLE 2

4,5-Dichloro-2-phenylpyrrole-3-carbonitrile and 5-chloro-2-phenylpyrrole-3-carbonitrile



To a magnetically stirred ice-water cooled solution of 2.00g (11.9 mmol.) of 2-phenyl-3-cyanopyrrole in 80 mL of methylene chloride is added dropwise over a period of 5 min., 1.90 mL (3.19 g, 23.6 mmol.) of sulfuryl chloride by means of a syringe. Throughout the addition the temperature is kept between 5 °C and 10 °C. Stirring at 5-10 °C is continued for 90 minutes. The reaction mixture is vacuum filtered to remove a precipitated solid (1.28g) identified as 5-chloro-2-phenylpyrrole-3-carbonitrile, mp 192.5-195 °C. The filtrate is diluted with 400 mL of ethyl acetate, washed twice with 200 mL of water, dried (sodium sulfate), treated with charcoal, filtered, and then concentrated in vacuo to give (after slurrying of the residue with hexane) 0.60g (21.3% yield) of a pink-purple solid. This solid is recrystallized from 5 mL of hot acetone to give 0.32g (9% yield) of 4,5-dichloro-2-phenylpyrrole-3-carbonitrile as an orangish brown solid, mp 254-255 °C. Max(mull,Nujol): 3165(br s), 3120(s), 2245(s), 1570(m), 1513(m), 1440(s), 1252(m), 1069(m), 996(m), 920-

(m), 768(s), 698(s), 665(s) cm^{-1} .

$^1\text{H-NMR}$ (DMSO): δ 7.73 (d, $J=6.6\text{Hz}$, 1.97H, two phenyl protons at C-2,6), δ 7.52 (t, $J=7.3\text{Hz}$, 2.04H, two phenyl protons at C-3,5), δ 7.44 (t, $J=7.3\text{Hz}$, 1.02H, one phenyl proton at C-4).


$^{13}\text{C-NMR}$ (DMSO): δ 137.51 (C-2 pyrrole carbon), δ 129.25 (C-4 phenyl carbon), δ 129.04 (C-3,5 phenyl carbons), δ 128.37 (C-1 phenyl carbon), δ 125.88 (C-2,6 phenyl carbons), δ 114.32 (either C-5 pyrrole or the nitrile carbon), δ 114.14 (either C-5 pyrrole or the nitrile carbon), δ 110.72 C-4 pyrrole carbon), δ 89.78 (C-3 pyrrole carbon).

Microanalysis (MW 237.09):

Calcd.:	C, 55.72%;	H, 2.55%;	N, 11.82%;	Cl, 29.91%
Found :	C, 55.78%;	H, 2.59%;	N, 11.12%;	Cl, 29.74%

EXAMPLE 3

p-Chloro- β -[(formylmethyl)amino]cinnamionitrile, diethyl acetal



A magnetically stirred solution of 250.00 g (1.39 mol.) of *p*-chlorobenzoylacetonitrile, 203 mL (185.95 g, 1.39 mol) of 2,2-diethoxyethylamine, and 1300 mL of dried toluene is heated at reflux for 20 hours. Water is collected in a Dean-Stark trap (23.8 mL, 95.2% theory). The hot cloudy dark brown solution with a large amount of undissolved solids is filtered through diatomaceous filter aid. After dilution with 200 mL of EtOAc, the solution is filtered through a 7cm X 13 5cm column of silica gel. The filtrate is concentrated *in vacuo* to give 354.38 g (86.4% crude yield) of a clear dark oil which slowly solidifies. This solid is recrystallised from hot cyclohexane to give 324.26g (79.1% yield) of a waxy orange solid. NMR of this product shows it to be composed of 78% (Z) and 23% (E) isomeric mixture of *p*-chloro- β -[(formylmethyl)amino] cinnamionitrile, diethyl acetal, m.p. 60-72° C. The following analytical data is for another similarly prepared sample.


Max(mull,Nujol): 3325(s), 3065(m), 2197(s), 1600(s), 1530(s), 1314(m), 1265(m), 1173(m), 1154(m), 1128(s), 1100(s), 1060(s), 1022(s), 939(m), 895(m), 844(s), 768(m), 730(m) cm^{-1} .

$^1\text{H-NMR}$ (chloroform): δ 7.47 (d, $J=8.6\text{Hz}$, 2.12H, two aromatic protons), δ 7.37 (d, $J=8.6\text{Hz}$, 2.12H, two aromatic protons), δ 5.10(E) & δ 4.86(Z) [br t, 1.25H, one N-H proton], δ 4.69(Z) & δ 4.60(E) [t, $J=5.1\text{Hz}$, 1.05H, one methine proton at the acetal carbon], δ 4.07 (E) & δ 4.05(Z) [s, 0.83H, enamine β proton], δ 3.71(E) & δ 3.68(Z) [q, $J=7.1\text{Hz}$, 2.22H, two methylene protons of one of two ethoxy groups], δ 3.56(Z) & δ 3.53(E) [q, $J=7.1\text{Hz}$, 2.22H, two methylene protons of one of two ethoxy groups], δ 3.18 (t, $J=5.1\text{Hz}$, 1.77H, two methylene protons of the ethyleneacetal group), δ 1.20 (t, $J=7.1\text{Hz}$, 4.90H, six methyl protons of the two ethoxy groups).

$^{13}\text{C-NMR}$ (chloroform): δ 161.21 (α -enamine carbon), δ 136.29 (Z) & δ 134.60(E) [either C-1 or C-4 of the phenyl ring], δ 134.08(Z) & δ 132.30(E) [either C-1 or C-4 of the phenyl ring], δ 129.34(Z) & δ 129.89(E) [either C-2,6 or C-3,5 of the phenyl ring], δ 128.94(Z) & δ 128.63(E) [either C-2,6 or C-3,5 of the phenyl ring], δ 121.19(Z) & δ 119.50(E) [nitrile carbon], δ 99.43(Z) & δ 100.63(E) [β -enamine carbon], δ 61.88(Z) & δ 63.25(E) [methine carbon of the acetal], δ 62.64(Z) & δ 63.03(E) [methylene carbons of the ethoxy groups], δ 46.32(Z) & δ 47.33(E) [methylene carbon of the ethyl amine group], δ 15.26 (methyl carbons of the ethoxy groups).

Microanalysis (MW 294.78):				
Calcd:	C, 61.11%;	H, 6.50%;	N, 9.51%;	Cl, 12.03%.
Found:	C, 61.25%;	H, 6.25%;	N, 9.34%;	Cl, 12.35%.

EXAMPLE 4

2(p-Chlorophenyl)-pyrrole-3-carbonitrile

To 108 mL of trifluoroacetic acid stirred at 23°C is added 54.00 g (0.183 mol) of solid p-chloro-β-[(formylmethyl)amino]cinnamionitrile, diethyl acetal over a period of 45 minutes. This addition produced an exotherm to 38°C and, 32 minutes into the addition, a solid started to precipitate. After stirring at room temperature for 30 minutes, the reaction mixture is vacuum filtered and the collected solid is washed first with trifluoroacetic acid, secondly with an ethyl acetate-hexane mixture, and finally with hexane. The yield is 16.83 g (45.4%) of an off-white solid, mp 165-166°C. The following anal. data is from a similarly prepared sample.


Max(mull, Nujol): 3275(br s), 2225(s), 1502(s), 1410(m), 1275(m), 1200(m), 1108(s), 1023(m), 999(m), 908-(m), 843(s), 752(s), 722(s), 695(s), 620(s) cm⁻¹.

H-NMR(acetone): δ11.22 (v br s, 0.99H, one pyrrole N-H proton), δ7.82 (d, J=8.9Hz, 2.46H, two aromatic phenyl protons), δ7.51 (d, J=8.9Hz, 2.46Hz, two aromatic phenyl protons), δ7.02 (t, J=2.6Hz, 1.01H, one pyrrole proton at C-5), δ6.58 (t, J=2.6Hz, 0.77H, one pyrrole proton at C-4).


C-NMR(acetone): δ137.73 (pyrrole C-2), δ134.42 (p-chlorophenyl at C-4), δ129.93 (methine carbons at C-3,5 of the phenyl ring), δ128.07 (methine carbons at C-2,6 of the phenyl ring), δ121.21 (pyrrole at C-5), δ117.93 (nitrile carbon), δ113.78 (pyrrole carbon at C-4), δ90.86 (pyrrole carbon at C-3).

Microanalysis (MW 202.64):				
Calcd.:	C, 65.19%;	H, 3.48%;	N, 13.83%;	Cl, 17.50%
Found:	C, 64.18%;	H, 3.52%;	N, 13.63%;	Cl, 17.74%

Use of the above procedure as shown or with the substitution of concentrated hydrochloric acid for trifluoroacetic acid affords the following compounds:



M and/or R	mp °C	Acid Used
4-Cl	165-166	conc. HCl, CF ₃ COOH
3,4-di-Cl	216-221	CF ₃ COOH
2-Cl	156-157	CF ₃ COOH
4-OCF ₃	143-145	CF ₃ COOH
4-CF ₃	179-180	CF ₃ COOH
2,4-di-Cl	197-199	CF ₃ COOH
3-Cl	150-156	CF ₃ COOH
4-CN	210-212	CF ₃ COOH
4-F	167-170	conc. HCl
4-SO ₂ CH ₃	221-221.5	CF ₃ COOH
3,4-di-F	173-175.5	CF ₃ COOH
3-CF ₃	166-168	CF ₃ COOH
4-COOCH ₃	155.5-158	CF ₃ COOH
4-CH ₃	117-137	CF ₃ COOH
4-NO ₂	174-177	CF ₃ COOH

EXAMPLE 5**4,5-Dichloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile**

To a mechanically stirred solution of 16.83g (83.1 mmol) of 2-(p-chlorophenyl)pyrrole-3-carbonitrile in 450 mL of glacial acetic acid at 36 °C is added dropwise 14.7 mL (24.70 g, 183.0 mmol) of sulfuryl chloride over a period of 18 minutes. The addition produces a slight exotherm to 39 °C and, after another 16 minutes, the reaction mixture is vacuum filtered. The collected solids are washed first with acetic acid and then with water. This solid after recrystallization from hot ethyl acetate, melts at 259-261 °C. By similar procedures other samples of this product were prepared and the analytical data for one such product is shown below.

Max(mull, Nujol): 3170(br s), 3100(m), 2225(s), 1508(m), 1097(m), 825(s), 717(m), 660(m) cm⁻¹.


H-NMR(DMSO): δ7.72 (d, J = 8.6Hz, 2.00H, two aromatic protons), δ7.56 (d, J = 8.6Hz, 2.00H, two aromatic protons).

C-NMR(DMSO): δ 136.01 (pyrrole C-2 carbon), δ 133.92 (p-chlorophenyl C-4 carbon), δ 129.09 (p-chlorophenyl C-3,5 carbons), δ 127.41 (p-chlorophenyl C-4 carbon), δ 127.11 (p-chlorophenyl C-1 carbon), δ 114.49 (nitrile carbon), δ 114.10 (pyrrole C-5 carbon), δ 110.92 (pyrrole C-4 carbon), δ 90.09 (pyrrole C-3 carbon).

Microanalysis (MW 271.54):				
Calcd.:	C, 48.65%;	H, 1.86%;	N, 10.32%;	Cl, 39.17%
Found:	C, 49.22%;	H, 2.12%;	N, 9.85%;	Cl, 39.03%

EXAMPLE 6


4,5-Dibromo-2-(α,α,α -trifluoro-p-tolyl)-pyrrole-3-carbonitrile



To a stirred mixture of 0.8g of 2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile in 70 mL of chloroform is added 2 mL of bromine. The mixture, on stirring overnight, deposits a white solid which is collected by filtration. Thin layer chromatography (1:1 ethyl acetate-hexane) shows a single component; m.p. $>230^\circ\text{C}$.

Anal. Calc'd for $\text{C}_{12}\text{H}_5\text{Br}_2\text{F}_3\text{N}_2$:				
	C, 36.55;	H, 1.27;	N, 7.11;	Br, 40.61.
Found:	C, 36.40;	H, 1.08;	N, 6.99;	Br, 40.55.

Following the procedures of Examples 5 and 6, but substituting the appropriately substituted phenylpyrrole-3-carbonitrile for 2-(α,α,α -trifluoro-p-tolyl) pyrrole-3-carbonitrile yields the following compounds.




<u>L</u>	<u>M</u>	<u>R</u>	<u>X</u>	<u>Y</u>	<u>mp °C</u>
H	H	4-NO ₂	Br	Br	274-277
H	H	4-F	Cl	Cl	>220
H	H	4-F	Br	Br	>220
H	H	4-SO ₂ CH ₃	Cl	Cl	>230
H	3-F	4-F	Cl	Cl	>230
H	3-F	4-F	Br	Br	>220
2-Cl	3-Cl	4-Cl	Cl	Cl	
2-Br	3-Br	4-Br	Br	Br	
H	H	4-OCF ₃	Cl	Cl	222-225
H	H	4-OCF ₃	Br	Br	231-232
H	H	4-OCF ₃	Cl	H	
H	H	4-CN	Br	Br	>230
H	H	4-CN	Cl	Cl	>240
H	H	4-SO ₂ CH ₃	Br	Br	>230
H	H	4-NO ₂	Cl	Cl	246-249
H	3-Cl	4-Cl	Br	Br	>260
H	H	3-CF ₃	Cl	Cl	>230
H	H	4-COCH ₃	Cl	Cl	251-254
H	2,3-CH=CH-		Cl	Cl	244-247
H	H	4-CH ₃	Cl	Cl	215-217
H	2-Cl	4-Cl	Br	Br	>230

	<u>L</u>	<u>M</u>	<u>R</u>	<u>X</u>	<u>Y</u>	<u>mp</u> ^{°C}
	H	H	3-Cl	Cl	Cl	>230
5	H	2-Cl	4-Cl	Cl	Cl	>230
	H	H	4-Cl	Br	Br	273-274
	H	H	2-Cl	Br	Br	>230
10	H	H	4-CF ₃	Cl	Cl	>230
	H	H	4-Br	Cl	Cl	>235
	H	H	2-Cl	Cl	Cl	>230
	H	3-Cl	4-Cl	Cl	Cl	>235
15	H	H	H	Cl	Cl	254-255
	H	H	4-Cl	Cl	Cl	255-257
	H	H	4-CF ₃	Br	Br	>230
20	H	H	4-Cl	Cl	Br	262-263 (dec.)
	H	H	4-Cl	Br	Cl	250-258 (dec.)
	H	3-Cl	5-Cl	Cl	Cl	>230
25	H	3-Cl	4-Cl	Cl	Br	>230
	2-Cl	4-Cl	5-F	Cl	Cl	207-210

30 EXAMPLE 73-Nitro-2-phenylpyrrole

35



40

Alpha-nitro acetophenone (5.7 g, 0.0345mol) is taken up in 100 mL toluene and 4.6g (0.0345mol) of amino acetaldehyde diethyl acetal is added. The reactants are put into a 250 mL RB flask fitted with a Dean-Stark trap. The trap is filled with 4A molecular sieves and the mixture is heated at reflux for 18 hours. The toluene is removed in vacuo to give 8.36 g of α -(2,2-diethoxyethylamino)- β -nitrostyrene as a brown oil. To this oil is added 50 mL of concentrated HCl. As the flask is swirled the oil turns to a yellow suspension. After 10 minutes the solid is filtered to give 2.48 g of a yellow solid. Recrystallization from ether/ethylacetate/hexane gives the product as two fractions, 2.08 g of m.p. 190-192 °C, (31%).

50 Max 1485 cm⁻¹(NO₂), H-NMR(CDCl₃/DMSO) δ 6.73(m,2H), 7.46(m,5H).

55


EXAMPLE 8

2,3-Dichloro-4-nitro-5-phenylpyrrole

5

10

15



20

A mixture of 3-nitro-2-phenylpyrrole (1.56g, 0.0083mol) in 60 mL of dioxane is cooled in an ice bath while 25.9g (.0182mol) of commercial sodium hypochlorite is added dropwise. After stirring for 45 minutes, the mixture is acidified with concentrated HCl. Water and Et₂O are added. The layers are separated and the top organic layer is washed with H₂O, dried over anhydrous MgSO₄ and concentrated in vacuo to give 2.21g of yellow solid. Purification by chromatography using silica gel and eluting with increasing ratios of ethyl acetate/ hexane gives, after stripping, 0.77g of yellow solid (36%) m.p. 190-190.5 °C;

25

Analysis: Calcd. for C ₁₀ H ₆ N ₂ O ₂ Cl ₂			
	C, 46.72;	H, 2.35;	N, 10.90
Found:	C, 46.96;	H, 2.86;	N, 10.02

30

Following the procedures of Examples 7 and 8 above but using the appropriately substituted α -nitroacetophenone and 2,2-di(C₁-C₄ alkoxy)ethylamine yields the substituted α -(2,2-di(C₁-C₄ alkoxy)-ethylamino)- β -nitrostyrene which is then converted to 3-nitro-2-(substituted)phenylpyrrole by treatment with HCl, HBr or CF₃CO₂H. Reaction of the thus formed substituted phenylpyrrole with sodium hypochlorite in dioxane yields the chloro analogs; whereas, reaction of the substituted phenylpyrrole with bromine in chloroform yields the bromine analogs.


35

40


45

50

55



L	M	R	X	Y	mp °C
H	H	H	Cl	Cl	190-190.5
H	4-Cl	H	Cl	Cl	214-215
H	4-Cl	H	Br	Br	203-204(dec.)
H	H	H	Br	Br	148.5-149
3-Cl	4-Cl	C	Cl	Cl	219-220(dec.)
H	4-Br	H	Cl	Cl	222-223(dec.)
H	H	4-CF ₃	Cl	Cl	166-168


EXAMPLE 94,5-Dichloro-2-(3,4-dichlorophenyl)-1-methyl-pyrrole-3-carbonitrile

In a 100 mL flask, 2 g of 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile in 60 mL dry THF gives a clear brown solution. 1 eq of KOtBu is added w/ stirring, this giving a clear solution after a few minutes. 1 eq of MeI is added by syringe and the solution is heated at reflux for 4 hours. It is then left to stir at RT overnight. The following day 50 mL of H₂O is added and the mixture extracted with 4 x 50 mL CHCl₃. The organic phases are combined, dried with MgSO₄, and concentrated. The resulting white solid is purified by flash chromatography on silica gel, using 50/50 EtOAc/hexane as an eluent. This gives 1.80 g of a white solid.

Yield = 86%

m.p. = 154-156 deg. C

Following the above procedure but substituting the appropriately substituted phenylpyrrole-3-carbonitrile or 3-nitro-2-(substituted)phenylpyrrole for 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile yields the compounds shown below.




<u>A</u>	<u>L</u>	<u>M</u>	<u>R</u>	<u>X</u>	<u>Y</u>	<u>mp</u> ^{°C}
CH ₃	H	H	4-Cl	Cl	Cl	152-153
C ₂ H ₅ OCH ₂	H	3-Cl	4-Cl	Cl	Cl	128-130
C ₂ H ₅	H	3-Cl	4-Cl	Cl	Cl	137-138
CH ₃	H	3-Cl	4-Cl	Cl	Cl	154-156
CH ₃	H	H	4-CF ₃	Br	Br	145-146
C ₆ H ₅ -CH ₂	H	H	4-CF ₃	Br	Br	145-147
C ₆ H ₅ -CH ₂	H	3-Cl	4-Cl	Cl	Cl	95-96
CH ₂ =CH-CH ₂	H	3-Cl	4-Cl	Cl	Cl	69-70
CH ₂ =C-CH ₂	H	3-Cl	4-Cl	Cl	Cl	
Cl						
CH=C-CH ₂	H	3-Cl	4-Cl	Cl	Cl	147-148
CH ₃ SCH ₂	H	3-Cl	4-Cl	Cl	Cl	
C(CH ₃) ₃	H	H	4-CF ₃	Cl	Cl	
CH ₃	H	H	4-CF ₃	Cl	Cl	99-100
CH ₃ SC ₂ H ₅	H	3-Cl	4-Cl	Cl	Cl	74-75
O						
C ₂ H ₅ -OC-CH ₂	H	3-Cl	4-Cl	Cl	Cl	118-120
C ₂ H ₅ -OCH ₂	H	H	4-CF ₃	Cl	Cl	99-100
CH ₃	H	H	4-OCH ₃	Br	Br	112-115
CH ₃	H	H	4-Cl	Br	Br	197-201
C ₂ H ₅ OCH ₂	H	H	4-OCF ₃	Cl	Cl	46-47
CH ₃	H	H	4-OCF ₃	Cl	Cl	72-73
C ₆ H ₅ -CH ₂	H	H	4-OCF ₃	Cl	Cl	oil
C ₂ H ₅ OCH ₂	H	H	4-Cl	Cl	Cl	-


	<u>A</u>	<u>L</u>	<u>M</u>	<u>R</u>	<u>X</u>	<u>Y</u>	<u>mp</u> ^{°C}
	HOCH ₂ CH ₂	H	3-Cl	4-Cl	Cl	Cl	143-145
5	NC	H	3-Cl	4-Cl	Cl	Cl	251-252
	C ₆ H ₅ CH ₂ OCH ₂	H	3-Cl	4-Cl	Cl	Cl	88-89
	Cl O-CH ₂	H	3-Cl	4-Cl	Cl	Cl	118-120
10	IC=C-CH ₂	H	3-Cl	4-Cl	Cl	Cl	115-116
	CH ₃	H	H	4-Cl	Br	CF ₃	126-129
	C ₂ H ₅ OCH ₂	H	H	4-Cl	Br	CF ₃	91-92
	C ₂ H ₅ -OCH ₂	H	3-Cl	4-Cl	Cl	Cl	118-120
15	C ₂ H ₅ -OCH ₂	H	H	4-Cl	Br	Br	104-105
	C ₆ H ₅ -CH ₂	H	H	4-Cl	Br	Br	81-82
	CH ₃	H	H	4-Cl	Br	Br	197-201
20	CN	H	H	4-CF ₃	Cl	Cl	138-139
	C ₂ H ₅ -OCH ₂	H	H	4-CF ₃	Br	CF ₃	104-105
	C ₂ H ₅ -OCH ₂	H	H	4-CF ₃	H	CF ₃	76-77
25	C ₂ H ₅ OCH ₂	H	3-Cl	4-Cl	Br	CF ₃	80-81

EXAMPLE 10

1-Benzyl-4,5-dibromo-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile




In a 100 mL flask, 1.5 g of 4,5-dibromo-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile is mixed with 50 mL dry THF to give a clear dark solution. 1 eq of KOtBu is added with stirring. After a few minutes the solution clears. Benzyl bromide (0.65 g) is added by syringe. The mixture is heated at reflux overnight. The following day TLC (50/50 EtOAc/hexane) indicates the presence of both starting material and product. The reaction is worked up in the following manner; 50 mL of water is added and the mixture is extracted with 4 x 50 mL CHCl₃. The organic phases are combined and washed with 4 x 50 mL 10% aq. NaOH. The organic phase is dried with MgSO₄ and stripped. This gives a brown solid which is crystallized from EtOAc/hexane. Yield = 0.75g = 40.7%
m.p. = 145-147 deg.C dec.

EXAMPLE 11**4,5-Dichloro-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)pyrrole-3-carbonitrile**

15 A sample of 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile (1.0g, 0.003 mole) is dissolved in 10 mL of dry tetrahydrofuran. To this solution is added potassium t-butoxide (0.37g, 0.0033 mole) followed by chloromethyl ethyl ether (0.312g, 0.0033 mole). The mixture is stirred for about 1 hour at room temperature and then poured into a large volume of water precipitating the product. The white solid is

20 collected and dried to give 1.0g (91%) with m.p. 128-130°.

EXAMPLE 12**4-Chloro-3-cyano-2-(p-chlorophenyl)pyrrole**

To a magnetically stirred 20°C solution of 17.87g (88.2 mmol, 1.00eq) of 2-(p-chlorophenyl)-3-cyanopyrrole in 800 mL of dioxane is added dropwise 250.15g (13.13g real, 176.4 mmol, 2.00eq) of 5.25

40 weight % bleach over a period of 30 minutes. After stirring at room temperature for a further 30 minutes, the reaction solution is poured into 2200 mL of water. The resulting mixture is vacuum filtered to remove a small amount of a black solid. The filtrate is acidified to pH 2 with concentrated HCl to produce a brown solid. This solid is vacuum filtered and the collected solids washed with water to give 22.41g of a brown solid. This solid is treated with 100 mL of 5% aqueous sodium hydroxide to dissolve the bulk of the material

45 while leaving a small amount of undissolved black solid. This black solid, dissolved into 100 mL of ethyl acetate, is washed with 75 mL each of 5% aqueous NaOH, water, and sat. aqueous NaCl. The ethyl acetate layer is dried (MgSO₄), treated with charcoal, filtered, and then rotary evaporated in vacuo to give 1.10g (5.3% yield) of an orangish brown solid. This solid is recrystallized from an ethyl acetate chloroform mixture to give 0.51g (2.4% yield) of an off-white solid of 4-chloro-3-cyano-2-(p-chlorophenyl)pyrrole. mp 251-


50 253.5°C.

EXAMPLE 13Preparation of 5-bromo-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile

5

10

15



20

A sample of 2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile (2.0g., 0.008 mole) is dissolved in 100 mL of dioxane by warming to 40,50°. Then the solution is cooled to 30°C and bromine (1.3g, .008 mole) is added. After stirring 1 hour at room temperature the solution is poured into water and a gray solid (2.2g, 88%) is collected. The mp is 233-236°C, decomposition.

In a similar fashion one can prepare 5-bromo-2-(3,4-dichloro)-3-nitropyrrole starting with 2-(3,4-dichlorophenyl)-3-nitropyrrole.


EXAMPLE 14

25

Preparation of 5-bromo-4-chloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile

30

35



40

A sample of 5-bromo-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile (0.158g, 0.005 mole) is dissolved in tetrahydrofuran (5 mL). An equivalent amount of t-butyl hypochlorite is added and the solution stirred overnight. The solution is poured into water and the precipitate (0.052g, 30%) is collected. The mp is >275°C.


In a similar fashion one can prepare 2-bromo-3-chloro-5-(3,4-dichlorophenyl)-4-nitropyrrole by starting with 2-bromo-5-(3,4-dichlorophenyl)-4-nitropyrrole.

50

55

EXAMPLE 15


Preparation of 5-bromo-4-chloro-2-(p-chlorophenyl)-pyrrole-3-carbonitrile



To a magnetically stirred 22°C solution of 0.17g (0.67 mmol., 1.00 equivalent) of 4-chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile in 100 mL of chloroform is added dropwise over a period of 30 minutes, a solution of 0.20 mL (0.62g, 3.88 mmol., 5.79 equivalent) of bromine in 5 mL of chloroform. The addition produces no exotherm. After stirring at room temperature for 3 1/4 hours, the clear red reaction solution is evaporated *in vacuo* to give 0.28g of an off-white solid. This solid is slurried with a hexanemethylene chloride mixture to give on vacuum filtration 0.23g of an off-white fluffy solid. mp 262-263°C; dec.

EXAMPLE 16


Preparation of 5-chloro-4-bromo-2-(p-chlorophenyl)-pyrrole-3-carbonitrile



To a magnetically stirred 45°C solution of 1.00g (4.22 mmol., 1.00 equivalent) of 5-chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile in 300 mL of chloroform is added dropwise over a period of 30 minutes, a solution of 0.40 mL (1.24g, 7.76 mmol., 1.84 equivalent) of bromine in 25 mL of chloroform. The addition produces no exotherm and towards the end of the addition, a small amount of a solid starts to precipitate. After stirring at room temperature for 19 1/2 hours the reaction mixture is evaporated *in vacuo* to give 1.49g of an orangish white solid. This solid is slurried with a hexane-methylene chloride mixture to give on vacuum filtration 1.33g (100% yield) of a fluffy white solid. mp 250-258°C, dec.

EXAMPLE 17


Preparation of 5-chloro-2-(p-chlorophenyl)pyrrole-3-
carbonitrile



To a 35 °C magnetically stirred solution of 2.40g (11.8 mmol., 1.00 equivalent) of 2-(p-chlorophenyl)-
 20 pyrrole-3-carbonitrile, and 65 mL of glacial acetic acid is added dropwise by syringe 0.75 mL (1.26g, 9.34
 mmol., 0.79 equivalent) of sulfuryl chloride over a period of 5 minutes. Approximately 5 minutes after the
 completion of the addition, a solid precipitated out of the reaction solution. After stirring at room temperature
 for 45 minutes, the reaction mixture is filtered and the collected solid is washed well with cold acetic acid to
 25 give 2.08g (74% crude yield) of an off-white solid. This solid is recrystallised from 75 mL of hot acetic acid
 to give 1.63g (58% yield) of 97 wt% pure. Product mp 258.5-261 °C.

EXAMPLE 18

Preparation of 2-(3,4-dichlorophenyl)-1-methylpyrrole-3-carbonitrile



40 In a 100 mL flask, 2.0 g of 2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile is dissolved in 50 mL of dry THF
 and 1 equivalent of potassium t-butoxide is added. This gives a slightly cloudy solution. One equivalent of
 methyl iodide is then added to the mixture by pipette. This leads to a slight lightening of the colour. A
 drying tube is attached to the flask and it is left to stir at ambient temperature overnight.

45 The next morning there is a slight light-coloured precipitate in the flask. 50 mL of water is then added
 and the solution becomes clear before a solid precipitates out of the solution. This solid is filtered out of the
 solution and compared to the starting material by TLC (25% ethyl acetate/hexane). This indicates a new
 single spot which is faster moving than the starting material. It is dried in a vacuum oven at 50 deg. C
 overnight. The product yield is 1.31g or 62% yield and has a melting point of 140-142 °C.

50


55

EXAMPLE 19

Preparation of 4,5-dichloro-2-(3,4-dichlorophenyl)-1-methylpyrrole-3-carbonitrile

5

10



15

In a 50 mL round bottom flask, 0.5g of 2-(3, 4-dichlorophenyl)-1-methylpyrrole-3-carbonitrile is mixed with 35 mL of glacial acetic acid. The mixture is warmed slightly with a heat gun to dissolve all of the pyrrole.

To this clear solution is added 2 eq. of sulfuryl chloride by pipette. The solution is left to stir at room temperature for 12 hours.

20

After 12 hours the solution is poured into 50 mL of water, resulting in a white precipitate. This is filtered out and dried in a vacuum oven at 50 °C for 3 hours.

The resulting solid is identical by TLC, (25% ethyl acetate/ hexane), and infrared analysis to the product of Example 9. Product yield is 0.36 (56%).


25

EXAMPLE 20

Preparation of 4,5-Dichloro-2-(3,4-dichlorophenyl)-1-(2-hydroxyethyl)-pyrrole-2-carbonitrile


30

35



40

45



50


To a stirred mixture of 2.0 g (6.5 mmol) of 4,5-dichloro-2-(3,4-dichlorophenyl)-pyrrole-3-carbonitrile and 0.88 g (7.8 mmol) of potassium *tert*-butoxide heated at reflux in 50 mL of dioxane is added 0.98 g (7.8 mmol) of bromoethanol. The mixture is stirred at reflux for 12 hours, cooled, diluted with 50 mL of water, and extracted several times with chloroform. The combined chloroform extracts are dried over magnesium sulfate and concentrated *in vacuo* to leave a solid which, on warming and dissolving in ethyl acetate, deposits on cooling mostly starting pyrrole. Concentration of the mother liquor and recrystallization of the residual solid from 20% ethyl acetate in hexane gives 0.31 g of a white solid, mp 143-145 °C; IR 5077A.

55

Anal. Calc'd for $C_{16}H_{23}NO_4$;				
	C, 44.57;	H, 2.29;	N, 8.00;	Cl, 40.57.
Found: (Agm 33139):	C, 44.77;	H, 2.29;	N, 8.06;	Cl, 40.14.


EXAMPLE 21

Preparation of 4,5-dichloro-2-(3,4-dichlorophenyl) pyrrole-1,3-dicarbonitrile



Potassium *t*-butoxide (617 mg, 55 mmol) is added in portions to a solution of 3-cyano-4,5-dichloro 2-(3,4-dichlorophenyl)pyrrole (1.52 g, 5 mmol) in anhydrous THF (20 mL). After 30 minutes, a solution of cyanogen bromide (583 mg, 5.5 mmol) in THF (1 mL) is added. The reaction mixture is stored at room temperature overnight. The solvent is removed in a rotary evaporator. The residue is treated with water and extracted with ethyl acetate. The organic layer is washed with water and saturated sodium chloride and dried ($MgSO_4$). Evaporation and crystallization of the residue from ethyl acetate gives white crystals (1.07 g); mp 250.5-252.0 °C; IR (nujol) 2255, 2245 cm^{-1} (CN); ^{13}C NMR ($DMSO-d_6$) 102.7 (N-CN), 113.7 (3-CN); Mass spectrum 331.9 ($M+1$).

Anal. Calc'd for $C_{12}H_3Cl_4N_3$ (330.99);				
	C, 43.54;	H, 0.91;	N, 12.70;	Cl 42.85.
Found:	C, 43.62;	H, 0.93,	N, 12.63;	Cl 41.95.

EXAMPLE 22Preparation of 4,5-Dichloro-2-(3,4-dichlorophenyl)-1-(3-iodo-2-propynyl)-pyrrole-3-carbonitrile

25

To a stirred mixture of 1.91 g (5.5 mmol) of 4,5-dichloro-2-(3,4-dichlorophenyl)-1-(2-propynyl)-pyrrole-3-carbonitrile in 500 mL of methanol is added 69 mL of 10% aqueous sodium hydroxide and then 0.70 g (2.7 mmol) of iodine. The mixture is stirred for 12 hours and then acidified and diluted with 200 mL of water. The precipitated solids are collected and recrystallized from methanol to afford 0.51 g white crystals, m.p. 115-116 °C.


30

This reaction is also applicable to the conversion of any of the formula III, IV, V, VI or VII substituted N-alkynylarylpyrroles of the present invention to N-substituted 3-iodo-2-propynyl arylpyrroles of said invention.

EXAMPLE 23

35

Preparation of 2-(3,4-dichlorophenyl)-4,5-diiodopyrrole-3-carbonitrile



50

N-iodosuccinimide (5.7 g, .0254 mol,) is added slowly to a solution of 2-(3,4-dichlorophenyl)-pyrrole-3-carbonitrile (3.0 g, .0127 mol) in 100 mL of THF. The reaction is stirred several hours at 25 °C until thin layer chromatography (silica gel; 100:100:1-ether:petroleum ether:acetic acid) shows completion. The mixture is evaporated in vacuo to give a residue containing the pyrrole and succinimide. The crude solid is dissolved in 500 mL of ether and shaken with 5 x 400 mL of water to remove the succinimide. The ether is dried over

55

Na₂SO₄ and evaporated in vacuo to leave 2.0 g (32.3%) of a grey-brown solid with mp >230 ° (loses purple vapors).

EXAMPLE 24

Preparation of 2-phenyl-1-pyrroline-4-carbonitrile

5

10

15

A solution of acrylonitrile (0.65 mL; 0.01 mol) and N-(trimethylsilyl)methyl-S-methyl-benzenethioimide (2.4 g; 0.01 mol) in THF (100 mL) is cooled to -5°C in an ice-acetone bath. Under a nitrogen purge, a solution of tetrabutylammonium fluoride (1.0 mL of a 1 N solution in THF) and THF (20 mL) is added dropwise over 30 minutes. The solution is stirred another 30 minutes at -5°C , and then allowed to warm slowly to ambient. Stirring is continued another 18 hours, and then solvent is removed under reduced pressure. The residue is partitioned between ether/water and the water layer extracted with fresh ether. The combined organic layer is washed with water, then saturated sodium chloride. The solution is dried over MgSO_4 , and cooling the filtrate causes precipitation of an off-white solid (1.2 g; 70% theoretical yield) whose spectral characteristics are identical to the material described by Tsuge [J. Org. Chem. 52, 2523 (1987)].

25

Calcd. for $\text{C}_{11}\text{H}_{10}\text{N}_2$: C, 77.65; H, 5.88; N, 16.47.
 Found: C, 77.55; H, 5.83; N, 16.39. mp = $95-97^{\circ}\text{C}$.

30


EXAMPLE 25

Preparation of 2-phenyl-pyrrole-4-carbonitrile

35

40

45



Under a nitrogen purge 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (0.23 g; 0.001 mol) and 2-phenyl-1-pyrroline-4-carbonitrile (0.17 g; 0.001 mol) is dissolved in 1,2-dimethoxyethane (13 mL) to form a clear orange solution. Pyridine (0.08 mL; 0.001 mol) is added in a single portion, causing a slight exotherm (to ca. 28°C) and an immediate formation of a green/grey precipitate. The suspension is stirred at room temperature for 18 hours during which time much of the solvent evaporates. The brownish semi-solid residue is partitioned between ether and a half-saturated solution of sodium carbonate. The red-brown aqueous layer is extracted twice with ether and the combined ether layer is washed with fresh water, then saturated sodium chloride. After drying with MgSO_4 , solvent is removed under reduced pressure to obtain a white semi-solid. This material was recrystallized from ethylene dichloride (DARCO treatment) to yield lavender crystals (0.1 g).

50


The identical product is obtained directly in a single step by condensing α -chloroacrylonitrile and N-(trimethylsilyl)methyl-S-methyl-benzenethioimide using tetrabutylammonium fluoride catalysis (analogous

to the preparation of 2-phenyl-1-pyrroline-4-carbonitrile described previously).

Calcd. for $C_{11}H_8N_2$: C, 78.57; H, 4.76; N, 16.67.
 Found: C, 78.65; H, 4.70; N, 16.43. m.p. - 155-158°C.

EXAMPLE 26

Preparation of 2,4-dibromo-5-phenyl pyrrole-3-carbonitrile




Under a nitrogen purge, a solution of bromine (0.6 mL; 0.012 mol) in $CHCl_3$ (5 mL) is added dropwise over 20 minutes to a stirring solution of 2-phenyl-pyrrole-4-carbonitrile (0.84 g; 0.05 mol) in $CHCl_3$ (20 mL). The resulting solution is stirred 18 hours at room temperature, then solvent is removed under reduced pressure to obtain a solid which is recrystallized from $C_2H_4Cl_2$ (DARCO treatment), yielding the desired final product (0.6 g), m.p. = 239-242°C.

Calcd. for $C_{11}H_6Br_2N_2$: C, 40.49; H, 1.84; Br, 49.08;
 N, 8.59.
 Found: C, 39.88; H, 1.87; Br, 48.81; N, 8.48.

By the procedure described in Example 24, 25 and 26, 2,4-dibromo-5-(p-chlorophenyl)pyrrole-3-carbonitrile, m.p. 270-272°C (dec.) is also prepared.

EXAMPLE 27

3',4'-Dichloro-3-(1,3-dioxolan-2-yl)-propiophenone




To a rapidly stirring mixture of magnesium turnings (0.64 g, 26 mmol) in 10 mL of tetrahydrofuran at 25°C in a 100 mL three-neck round bottom flask equipped with a thermometer, a 60 mL addition funnel, and a nitrogen inlet is added dropwise 2-(2-bromoethyl)-1,3-dioxolane (4.7 g, 26 mmol) in 40 mL of tetrahydrofuran. The rate of addition is adjusted so as to maintain the reaction temperature below 50°C. The reaction is then allowed to stir for 1 hour at 25°C. 120 mL of tetrahydrofuran is mixed with potassium

3,4-dichlorobenzoate (5.0 g, 22 mmol) under a blanket of nitrogen. The Grignard solution is then quickly decanted away from the unreacted magnesium turnings, and added dropwise to the rapidly stirring potassium benzoate suspension. The reaction is then allowed to stir for 24 hours at 25 °C. Fifty mL of diethyl ether and 15 mL of 3N hydrochloric acid are added to the reaction mixture and the layers separated.

5 The organic layer is washed with saturated aqueous sodium bicarbonate until neutral followed by one washing with 10 mL of brine. Drying over sodium sulfate, and rotary evaporation yields a beige semisolid which is chromatographed over silica gel using 3:1 hexane-ethyl acetate as eluent to give the keto-acetal (4.3 g, 60%) as a white solid, m.p. 115-117 °C.

10 EXAMPLE 28

Preparation of 3-(3,4-dichlorobenzoyl)propionaldehyde




Ten grams (26 mmol) of 3',4'-dichloro-3-(1,3-dioxolan-2-yl)-propiophenone is added to 30 mL of 0.2M

25 oxalic acid (made by dissolving 0.9 g of oxalic acid dihydrate in 30 mL of water) and 5 mL of ethanol. The mixture is refluxed for 1 hour and then allowed to cool. Most of the ethanol is rotary evaporated off and 100 mL of diethyl ether is added along with 20 mL of saturated aqueous sodium bicarbonate. The layers are separated and the organic phase is dried over magnesium sulfate. Rotary evaporation yields a viscous yellow oil which is chromatographed over silica gel using 3:1 hexane-ethyl acetate to give the keto-aldehyde

30 (6.3 g, 75%) as a white solid.

EXAMPLE 29

Preparation of 2-(3,4-dichlorophenyl)pyrrole




To a suspension of 3-(3,4-dichlorobenzoyl) propionaldehyde (6 g, 26 mmol) in 60 mL of absolute ethanol is added ammonium acetate (4 g, 52 mmol). The reaction is refluxed for 20 minutes and allowed to cool. Most of the ethanol is rotary evaporated and 200 mL of 1:1 dichloromethane-diethyl ether along with

50 50 mL of water is added. The layers are separated and the organic phase is dried over sodium sulfate. Rotary evaporation yields a dark brown oil which is chromatographed over silica gel using 3:1 hexane-ethyl acetate as eluent to give the pyrrole (4.6 g, 83%) as a light brown solid, m.p. 49-51 °C.

55


EXAMPLE 30Preparation of 5-(3,4-dichlorophenyl)pyrrole-2-carboxaldehyde

5



15 To 10 mL of dimethylformamide stirring under nitrogen in a 50 mL round bottom flask is added phosphorus oxychloride (0.6 mL, 6.5 mmol) dropwise via syringe. The solution, warms and becomes light yellow in color. It is allowed to stir for 20 minutes before the portionwise addition of 2-(3,4-dichlorophenyl)-pyrrole (1 g, 4.7 mmol). The beige suspension which results is allowed to stir for 30 minutes before being heated to 50 °C for 40 minutes. A solution of sodium acetate (10 g, 122 mmol) in 15 mL of water is added

20 to the cooled reaction which is then allowed to stir for 20 minutes. A beige precipitate is filtered off from the reaction mixture and air-dried for 20 hours to give the essentially pure aldehyde (1.1 g, 95%), mp > 200 °C.


EXAMPLE 31Preparation of 5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile

35 To a suspension of 5-(3,4-dichlorophenyl)pyrrole-2-carboxaldehyde (1.5 g, 6.2 mmol) in 20 mL of water and 20 mL of ethanol, is added hydroxylamine-O-sulfonic acid (0.7 g, 6.2 mmol). The reaction is refluxed for 1 hour during which time a gray precipitate appears. After being allowed to cool, the reaction is filtered to give essentially pure nitrile (1.5 g, 99%) as a gray solid, m.p. 170-171 °C.

40

EXAMPLE 32Preparation of 3,4-dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile

45




55

To a solution of 5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile (0.5 g, 2.1 mmol) in 20 mL of tetrahydrofuran under nitrogen is added portionwise N-bromo-succinimide (0.8 g, 4.2 mmol). The reaction is stirred at 25 °C for 30 minutes before the addition of 10 mL of water and 40 mL of diethyl ether. The layers

are separated and the organic layer dried over sodium sulfate. Rotary evaporation is followed by chromatography over silica gel using 3:1 hexane-ethyl acetate as eluent to afford the dibromopyrrole (0.5 g, 60%) as a brown solid, m.p. > 250 °C.

5 EXAMPLE 33

Preparation of 4-phenylpyrrole-3-carbonitrile




25 To a mixture of 5.0 g (39 mmol) of cinnamionitrile and 7.6 g (39 mmol) of (p-tolylsulfonyl)methyl isocyanide in 35 mL of DMSO and 65 mL of ether is added over a 20 minute period a suspension of 1.86 g of a 60% oil suspension of sodium hydride (1.11 g; 46 mmol) in 80 mL of ether. The reaction mixture is maintained under nitrogen for an hour and then diluted with ether and water. The ether layer is separated, dried over magnesium sulfate, and concentrated in vacuo. The resulting oil is chromatographed on silica gel using 1:1 chloroform ethyl acetate to give 2.5 g of cream-colored solids. Recrystallization from ether-hexane affords 1.15 g, m.p. 123-125 °C; NMR M86-1077.

30 Lit.: Tet. Letters 5337 (1972): m.p. 128-129 °C.

EXAMPLE 34


35 Preparation of 2,5-dichloro-4-phenylpyrrole-3-carbonitrile



To a stirred mixture of 0.66 g (3.9 mmol) of 4-phenylpyrrole-3-carbonitrile in 20 mL of dry THF cooled to 6 °C with an ice-water bath is added from a syringe 0.66 mL (1.11 g; 8.2 mmol) of sulfuryl chloride over a 4 minute period. The mixture is maintained at 5-10 °C for an additional 45 minutes and then stirred an additional 30 minutes with the ice bath removed. After the reaction mixture is poured into 80 mL of ethyl acetate and 40 mL of water, the organic phase is separated, washed with water, and dried over sodium sulfate. Filtration through a short column of silica gel, rinsing with ethyl acetate, and concentration of the combined filtrate in vacuo gives 0.95 g of dark solid. Recrystallization from chloroform gives 0.42 g of off-white crystals, m.p. 195-196 °C (dec.).

Anal. Calcd for C ₁₁ H ₆ Cl ₂ N ₂ :				
	C, 55.72;	H, 2.55;	N, 11.82;	Cl, 29.91.
Found:	C, 55.66;	H, 2.65;	N, 11.69;	Cl, 29.97.


Following the procedures of Examples 33 and 34, the following analogs are prepared. For the synthesis of 2,6-dibromo-4-(p-chlorophenyl)pyrrole-3-carbonitrile, the procedure of Example 33 is followed using bromine in dioxane to replace sulfonyl chloride and tetrahydrofuron.



R	X	Y	m.p. °C
4-Cl	Cl	Cl	237-240 (dec.)
4-CH ₃	Cl	Cl	103-206
4-Cl	Br	Br	> 245 °

EXAMPLE 35

Ethyl 4-(p-chlorophenyl)-pyrrole-3-carboxylate



To a mixture of 5.63 g of a 60% sodium hydride/oil suspension in 200 mL of dry ether under nitrogen is added from an additional funnel a mixture of 23.5 g (122 mmol) of ethyl p-chlorocinnamate and 19.4 g (122 mmol) of (p-tolylsulfonyl)methyl isocyanide in solution in 180 mL of ether and 80 mL of dimethylsulfonide. The addition time is about 20 minutes and results in gentle refluxing of the mixture. After another 10 minutes stirring, the mixture is diluted with 100 mL of water. The mixture is extracted four times with ether which is then dried over magnesium sulfate followed by concentrated in vacuo. The resulting solid is

recrystallized from ethylene dichloride to give 7.8 g of crystals, m.p. 137-138 °C.


Anal. Calcd for $C_{13}H_{12}ClNO_2$:

	C, 62.53;	H, 4.81;	N, 5.61;	Cl, 14.23.
Found:	C, 61.31,	H, 5.12;	N, 5.32;	Cl, 14.57.

Concentration of the mother liquor for the crystallization leaves additional crude ester which is carried on to the saponification step.

EXAMPLE 36

Preparation of 3-(p-chlorophenyl)-pyrrole



A mixture of 22.0 g of crude ethyl 4-(p-chlorophenyl)-pyrrole-3-carboxylate from the recrystallization mother liquor and the recrystallized product from the previous step is stirred at reflux with 150 mL of 10% aqueous sodium hydroxide for 2.5 hours. The mixture is cooled, extracted with ether, and acidified to give a precipitate which on collection and drying weighs 11.6 g.

A mixture of 10.5 g of the acid in 100 mL of β -ethanolamine is heated at reflux for three hours. After cooling, the mixture is poured over 400 mL of ice and the resulting mixture is extracted four times with chloroform. The chloroform solution, after drying over magnesium sulfate and treatment with activated charcoal, is concentrated *in vacuo* to leave a brown solid. Chromatography on silica gel using 1:1 ethyl acetate hexane gives 4.0 g of a white solid, m.p. 117-118 °C.

EXAMPLE 37Preparation of 3-(p-chlorophenyl)-pyrrole-2-carboxal-
dehyde


5

10

15

20

25



To a mixture of 0.86 g (12 mmol) of dimethylformamide in 10 mL of ethylene dichloride maintained under nitrogen and cooled in an ice bath is added 1.49 g (12 mmol) of oxalyl chloride in 10 mL of ethylene dichloride over a period of 25 minutes. The ice bath is removed, the mixture is stirred an additional 15 minutes and recooled in an ice bath. To this mixture is added 1.5 g (8.5 mmol) of 3-(p-chlorophenyl)-pyrrole in 25 mL of ethylene dichloride over a 20 minute period. The ice bath is removed and after an additional 30 minutes of stirring, the mixture is poured into 50 mL of ice-water and 6 mL of 50% sodium hydroxide. The resulting mixture is extracted with ether and with chloroform and the combined organic mixture is dried over magnesium sulfate and concentrated in vacuo. Purification of the resulting solid by chromatography on silica gel using 1:1 ethyl acetate hexane gives 0.63 g of off-white solid which is used directly for conversion to 3-(p-chlorophenyl)-pyrrole-2-carbonitrile.

35

EXAMPLE 38


40

Preparation of 3-(p-chlorophenyl)-pyrrole-2-carboni-
trile

45

50

55



A mixture of 0.63 g (3.1 mmol) of 3-(p-chlorophenyl)-pyrrole-2-carboxaldehyde in 10 mL of water is stirred and ice-cooled while 0.52 g (4.6 mmol) of hydroxylamine-O-sulfonic acid in 10 mL of water is slowly added. After the addition, the cooling bath is removed and the mixture is heated for 25 minutes. On cooling, the resulting solid is collected and shown, by NMR, to be a mixture of product and starting aldehyde. This mixture is reacted in the same manner with an additional 0.49 g (4.2 mmol) of hydroxylamine-O-sulfonic acid in a total of 30 mL of water. The mixture is heated at 60-70 °C for 2 hours. The mixture is cooled and the resulting solids are collected and purified by chromatography or silica gel using 1:1 ethyl acetate hexane to give 0.40 g of pink solid, m.p. 114-115 °C.


10 EXAMPLE 39

Preparation of 4,5-Dibromo-3-(p-chlorophenyl)-pyrrole-2-carbonitrile

15

20

25



30

To a mixture 0.40 g (2.0 mmol) of 3-(p-chlorophenyl)pyrrole-2-carbonitrile in 25 mL of chloroform is added 0.63 g (4.0 mmol) of bromine. After 20 minutes, the precipitate which forms is collected and recrystallized from ethyl acetate to give 0.21 g of pink crystals, m.p. > 250 °C.

35

Anal. Calcd for $\text{C}_{11}\text{H}_5\text{Br}_2\text{ClN}$:					
	C, 36.62;	H, 1.39;	Br, 44.38;	Cl, 9.85;	N, 7.77.
Found:	C, 36.92;	H, 1.32;	Br, 44.62;	Cl, 9.88;	N, 7.50


40

45

50

55


EXAMPLE 40

Preparation of Ethyl 5-bromo-4-(p-chlorophenyl)pyrrole-3-carboxylate


30 Ethyl 4-(p-chlorophenyl)pyrrole-3-carboxylate (1.6 g., 0.0064 mmol) is dissolved in tetrahydrofuran (40 mL). N-bromosuccinimide (1.14 g., 0.0064 mmol) is added in small portions at 25-28°C. After the addition is complete, the solution is stirred overnight at room temperature. The solution is concentrated in vacuo and the solid residue partitioned between water and ether. The ether layer is separated and dried over magnesium sulfate. Work-up of the ether extract leaves 1.9 g (90%) of a white solid which is purified by stirring with a mixture of 80/20 hexane/ethyl acetate. The insoluble solid (1.3 g, 62%) is collected and has
35 m.p. 161-164°C.

Calcd for C ₁₃ H ₁₁ BrClNO ₂ :					
Found:	C, 47.50; C, 47.39;	H, 3.34; H, 3.38;	N, 4.26; N, 4.12;	Br, 24.33; Br, 24.29;	Cl, 10.80. Cl, 10.77


EXAMPLE 41

Preparation of 5-bromo-4-(p-chlorophenyl)pyrrole-3-carboxylic acid

Ethyl 5-bromo-4-(p-chlorophenyl)pyrrole-3-carboxylate (15 g., 0.045 mmol) is added to 200 mL of 10% sodium hydroxide and the slurry heated to reflux. After everything appears to dissolve the mixture is refluxed an additional 40 minutes. The mixture is cooled, filtered and the filtrate acidified. The white precipitate (8.0 g, 58%) is collected and dried. The solid has m.p. >205°C and an NMR (d_6 -DMSO) which showed a pyrrole proton at 7.52 (d). The mass spectrum is also consistent for a monobrominated compound.

EXAMPLE 42**Preparation of 2-bromo-3-(p-chlorophenyl)pyrrole**

5-bromo-4-(p-chlorophenyl)pyrrole-3-carboxylic acid (8.0 g., 0.026 mmol) is added to aminoethanol (24 mL) and the slurry slowly warmed to 110-120°C and held at that temperature for 1 hour. The solution is cooled and poured into water and extracted with ether. The ether extract, by thin layer chromatography (75/25, hexane/ethyl acetate), shows a major fast moving spot and a slower moving minor component. Work-up of the ether leaves a dark solid (4.0 g., 56%) which is 2-bromo-3-(p-chlorophenyl)pyrrole and is used immediately to prepare 5-bromo-4-(p-chlorophenyl)pyrrole-2-carbonitrile.


EXAMPLE 43**Preparation of 5-bromo-4-(p-chlorophenyl)pyrrole-2-carbonitrile**

A freshly prepared sample of 2-bromo-3-(p-chlorophenyl)pyrrole (4.0 g., 0.015 mmol) is dissolved in dry dimethoxyethane (25 mL). Then while holding the temperature below 25°C, chlorosulfonyl isocyanate (3.08 g., 0.022 mmol) is added. After stirring overnight, the solution is treated with dimethylformamide (6 mL) and stirred for 3 hours. Finally, the solution is poured into water precipitating a brown solid (3.8 g, 90%). Dry column chromatography (80/20 hexane/ethyl acetate) yields 1.4 g (33%) of white solid with m.p. 202-204°C.

Calcd for $C_{11}H_6BrC_1N_2$:					
	C, 46.90;	H, 2.13;	N, 9.95;	C1, 12.61;	Br, 28.39.
Found:	C, 47.20;	H, 2.09;	N, 9.80;	C1, 12.36;	Br, 27.42.

EXAMPLE 44

Preparation of 3,5-Dibromo-4-(p-chlorophenyl)pyrrole-2-carbonitrile




A sample of 5-bromo-4-(p-chlorophenyl)pyrrole-2-carbonitrile (2.2 g., 0.0078 mol) is dissolved in 30 mL of dry dioxane. The solution is heated with bromine (1.3 g., 0.008 mol) in dioxane (20 mL) and then stirred overnight at room temperature. The reaction mixture is poured into water precipitating a tan solid (2.6 g., 92%). A portion (1.6 g) is purified by flash chromatography using 75/25 hexane/ethyl acetate to give 0.8 g of grey solid with m.p. 191-194 °C.

Calcd for C ₁₁ H ₅ Br ₂ CN:					
	C, 36.61;	H, 1.38;	N, 7.76;	Cl, 9.84;	Br, 44.3.
Found:	C, 37.46;	H, 1.25;	N, 7.41;	Cl, 9.53;	Br, 42.99.

EXAMPLE 45

Preparation of 3-(3,4-dichlorophenyl)-4-nitropyrrole




Sodium hydride (2.66 g of a 60% suspension in oil is rinsed with dry ether; 66 mmol) and suspended in 150 mL of dry ether. To this mixture is added over 15 minutes a mixture of 12.0 g (5.5 mmol) of 3,4-dichloro- β -nitrostyrene and 10.8 g (5.5 mmol) of (p-tolylsulfonyl)methyl isocyanide in 50 mL of DMSO and 150 mL of ether. The mixture is stirred for 1.5 hours and then diluted with 150-200 mL of water and additional ether. The ether layer is separated, dried over magnesium sulfate, and concentrated in vacuo.

The resulting 10.6 g of crude product is purified by chromatography on silica gel using a 4:1 mixture of chloroform and ethyl acetate. A 7.2 g solid fraction is recrystallized from chloroform-ethyl acetate-hexane to give 3.0 g of yellow solid, m.p. 187-188 °C (dec.).

Anal. Calcd for C ₁₀ H ₆ Cl ₂ N ₂ O ₂ :			
	C, 46.72;	H, 2.35;	N, 10.90.
Found:	C, 46.96;	H, 2.60;	N, 9.77

EXAMPLE 46


Preparation of 2,5-Dichloro-3-(3,4-dichlorophenyl)-4-nitropyrrole



To a mixture of 3-(3,4-dichlorophenyl)-4-nitropyrrole (2.5 g, 9.7 mmol) warmed to about 40 °C in 200 mL of chloroform is added over one minute 2.95 g (22 mmol) of sulfuryl chloride. After another hour, the mixture is diluted with 100 mL of saturated sodium bicarbonate solution and 300 mL of ether. The organic layer is separated and dried over magnesium sulfate. Concentration, *in vacuo*, leaves a brown solid which is chromatographed on silica gel using 4:1 chloroform ethyl acetate. An orange solid fraction is recrystallized from chloroform and then rechromatographed on silica gel using 4:1 chloroform ethyl acetate to yield 0.36 g of yellow solid, m.p. 193-194 °C.

Also prepared by procedure of Examples 45 and 46 above is 2,5-dichloro-3-nitro-4-phenylpyrrole, m.p. 193-194 °C(dec.).

EXAMPLE 47

Preparation of 5-(p-chlorophenyl)pyrrole-2,4-dicarbo-
nitrile


30

A sample of 2-p-chlorophenyl-3-cyanopyrrole, prepared by the method of Example 4, (3.0 g, 0.015 mole) is dissolved in 50 mL of dry dimethoxyethane. To this solution is added chlorosulfonyl isocyanate (3.39 g, 0.024 mole). The addition is exothermic and some cooling is necessary. After stirring 3 hours at room temperature, dimethylformamide (6-7 mL) is added and the solution is stirred 4 hours more. The

35

solution is then poured into water precipitating a white solid (3.4 g, 100%). A sample (1.0 g) is purified by dissolving in ethyl acetate and then passing the solution through a 60 mL coarse filter funnel packed with silica gel. The filtrate is concentrated to yield 0.7 g of a white solid with m.p. 235-240 °C.


Following the procedure of Example 47, the following analogs are prepared:



R	L	m.p.
3-C1	4-C1	>225 °C
H	4-OCF ₃	185-190 °C
H	4-CF ₃	180-185 °C


EXAMPLE 48

Preparation of 3-Bromo-5-(p-chlorophenyl)pyrrole-2,4-dicarbonitrile



A sample of 5-(p-chlorophenyl)pyrrole-2,4-dicarbonitrile (1.0 g, 0.004 mole) is dissolved in 20 mL of dioxane and a solution of bromine (0.8 g, 0.005 mole) in dioxane (10 mL) is then added thereto. The solution is stirred several hours at room temperature and then poured into water precipitating a white solid (1.2 g, 100%). The solid has a m.p. >225 °C and a mass spectrum of a sample gives a pattern consistent with the desired structure.


Following the procedure set forth above in Example 48, the following additional compounds are prepared:



R	L	m.p.
3-C1	4-C1	>250 °C
H	4-OCF ₃	218-223 °C
H	4-CF ₃	239-241 °C

EXAMPLE 49

Preparation of bromofumaronitrile



15 Under a nitrogen purge, fumaronitrile (15.6 g; 0.2 mol) in CHCl_3 (150 mL) is heated to reflux, resulting in a clear solution. A solution of bromine (5.3 mL; 0.2 mol) in CHCl_3 (25 mL) is added dropwise over 30 minutes, resulting in a slow decolorization and acidic (pH test paper) fumes being released. The solution is refluxed another 90 minutes, during which time most of the color has been discharged. The solution is cooled and solvent is removed under reduced pressure, leaving an amber oil (weight approximately theoretical for bromofumaronitrile). The oil is subjected to bulb-to-bulb distillation (0.2 mm Hg), maintaining the temperature below 120°C (above that point, a rapid decomposition of material occurs). A semi-solid is obtained which slowly forms a waxy, amber solid, m.p. -43 – 47°C .

25


Calcd for C_4HBrN :			
	C, 30.57;	H, 0.64;	N, 17.83.
Found:	C, 29.13;	H, 0.75;	N, 16.94.

30

EXAMPLE 50

Preparation of 2-phenyl-pyrrole-3,4-dicarbonitrile

35



Under a nitrogen purge, a solution of bromofumaronitrile (4.7 g; 0.03 mol) and N-(trimethylsilyl) methyl-S-methyl-benzene-thioimide (7.1 g; 0.03 mol) in hexamethylphosphoramide (HMPA) (35 mL) is stirred at


room temperature. In a single portion, water (1.6 mL; 0.09 mol) is added, washed in with HMPA (10 mL). The solution almost immediately begins to exotherm, the temperature rapidly reaching 100 °C before subsiding. The resulting dark red solution is allowed to stir at ambient temperature 20 hours. Pouring the reaction mixture onto an ice/water mixture results in a gummy material which slowly yields a discreet beige solid. This material is collected by filtration and washed with cold water and dried on the filter. After further drying (vacuum oven; 60 °C), the material is twice recrystallized from C₂H₄Cl₂ (DARCO treatment) to yield a white powder.

Calcd for C ₁₂ H ₇ N ₃ :			
	C, 74.61;	H, 3.63;	N, 21.76.
Found:	C, 74.45;	H, 3.84;	N, 21.61.

m.p. = 197-200 °C.

EXAMPLE 51


Preparation of 2-bromo-5-phenylpyrrole-3,4-dicarbonitrile



Under a nitrogen purge, 2-phenyl-pyrrole-3,4-dicarbonitrile (1.4 g; 0.0075 mol) is added to CHCl₃ (35 mL), much of the solid dissolving. A solution of bromine (0.4 mL; 0.008 mol) in CHCl₃ (5 mL) is added dropwise over 20 minutes. Initially the color is discharged rapidly, but as a new, gummy solid begins to precipitate, the color remains. After stirring 30 minutes at ambient, the mixture is brought to reflux, resulting in a much more discreet solid. After refluxing 90 minutes, the reaction mixture is cooled and an aliquot is removed and analyzed (HPLC), showing ca. 60% starting material still remaining. In a single portion fresh bromine (0.2 mL; 0.004 mol) is added, and refluxing continued another 45 minutes whereupon an aliquot shows 10% starting material remaining. Another fresh portion of bromine (0.2 mL; 0.004 mol) is added to the refluxing suspension and refluxing is continued another 30 minutes. The suspension is cooled and stirred 18 hours at room temperature. Solvent is removed under reduced pressure to yield a greenish solid which is extracted with hot CHCl₃, leaving behind a dark residue. The extract is treated with DARCO and filtered hot. The clear yellow filtrate quickly began to deposit a white precipitate. After cooling to -10 °C, the white solid is collected by filtration.

Calcd for C ₁₂ H ₅ BrN ₃ :				
	C, 52.94;	H, 2.21;	N, 15.44;	Br, 29.41.
Found:	C, 51.64;	H, 2.35;	N, 14.91;	Br, 28.69.

m.p. = 225-258 °C.


EXAMPLE 52Preparation of 2-(3,4-Dichlorophenyl)-5-nitropyrrole-3-carbonitrile

2-(3,4-Dichlorophenyl)pyrrole-3-carbonitrile (3.0 g, 0.013 mole) is added to acetic anhydride (50 mL) and 90% nitric acid (0.6 ml) with very little exotherm. The mixture is slowly warmed to 30° and is then held at 30-33° until everything goes into solution. Gradually a new solid precipitates. The mixture is stirred for 2 to 3 hours at room temperature and then poured into water and ice to decompose the acetic anhydride. After stirring 1 hour the mixture is filtered and the solid (2.9 g, 82%) collected and dried. A portion (1.5 g) is purified by column chromatography on silica gel using 75/25 hexane/ethyl acetate for elution to give 0.7 g of yellow solid with m.p. 228-231°.

25

Calcd for C ₁₁ H ₅ Cl ₂ N ₃ O ₂ :				
	C, 46.80;	H, 1.77;	N, 14.89;	Cl, 25.17
Found:	C, 46.50;	N, 1.96;	N, 14.27;	Cl, 24.30.

30 By the same procedure, starting with 2-(p-chlorophenyl)pyrrole-3-carbonitrile, 2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile is obtained, m.p. 201-206° C. Also, 2-(p-trifluoromethylphenyl)pyrrole-3-carbonitrile gives 2-(p-trifluoromethylphenyl)-5-nitropyrrole-3-carbonitrile by the above procedure. This compound has a melting point of 164-165.5° C.

EXAMPLE 53Preparation of 4-Bromo-2-(3,4-dichlorophenyl)-5-nitropyrrole-3-carbonitrile


55 2-(3,4-Dichlorophenyl)-5-nitropyrrole-3-carbonitrile (0.5 g, 0.0017 mol) is dissolved in dry dioxane (10 mL). To this solution is added bromine (0.28 g, 0.0017 mole) in dioxane. After stirring overnight, the solution is poured into water precipitating a tan solid (0.54 g, 88%). Recrystallization from acetonitrile (5 mL) gives 0.26 g of tan solid with m.p. 195-200° C.

Calcd for C ₁₁ H ₄ BrCl ₂ N ₃ O ₂ :					
	C, 36.57;	H, 1.10;	N, 11.63;	Br, 22.13;	Cl, 19.67.
Found:	C, 36.46;	H, 1.29;	N, 11.50;	Br, 21.63;	Cl, 19.28.

Following the above procedure of Example 53, but starting with 2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile gives 4-bromo-2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile, m.p. 180-185 °C.

EXAMPLE 54


5-(3,4-Dichlorophenyl)-4-nitropyrrole-2-carbonitrile



To a suspension of 5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile (1.2 g, 5.1 mmol) in 25 mL of acetic anhydride at 30 °C under nitrogen, is added dropwise 90% nitric acid (0.3 mL, 5.1 mmol). The reaction exotherms to 45 °C and becomes a green solution. After being allowed to stir for 2 hours the reaction is poured into 50 mL of water and stirred vigorously for 5 minutes. The beige precipitate which results is filtered off and dissolved in a minimum amount of acetone. Chromatography over silica gel using 3:1 hexane-ethyl acetate affords the nitropyrrole (1.2 g, 84%) as an off-white solid, m.p. >200 °C.

EXAMPLE 55

3-Bromo-5-(3',4'-dichlorophenyl)-4-nitropyrrole-2-carbonitrile



To a suspension of 5-(3,4-dichlorophenyl)-4-nitropyrrole-2-carbonitrile (0.6 g, 2.1 mmol) in 10 mL of dioxane at 25 °C, under nitrogen, is added dropwise a solution of bromine (0.3 g, 2.1 mmol) in 5 mL of dioxane. The reaction is allowed to stir overnight. Addition of 50 mL of water causes precipitation of a yellow solid which is collected and vacuum oven dried (50 mm Hg, 45 °C) to afford the brominated pyrrole (0.7 g, 90%) as a light yellow solid, m.p. >200 °C.

EXAMPLE 56

4-(p-chlorophenyl)-2-(trifluoromethyl-2-oxazolin-5-one

In a single portion, trifluoroacetic anhydride, (1.7 mL; 0.012 mol) is added to powdered 2-(p-chlorophenyl)glycine (11.4 g; 0.06 mol), causing an immediate exotherm to about 40 °C, a yellow color


forming on the surface of the solid. As the mixture is slowly heated to 70 °C, more of the solid dissolves to an orange/amber oil. All the solid dissolved in approximately 2 hours, and heating is continued another hour. Solvent is removed under reduced pressure on a rotary evaporator. Toluene is twice added and removed under reduced pressure, but the odor of trifluoroacetic acid is still evident. This yellow semi-solid (yield theoretical; purity > 90% by HPLC) is the above-identified compound and is used in the next step without further purification.

EXAMPLE 57

Preparation of 2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile

4-(p-chlorophenyl)-2-(trifluoromethyl)-2-oxazolin-5-one (2.5 g; 0.01 mol) is dissolved in nitromethane (50 mL). In a single portion, 2-chloroacrylonitrile (8.0 mL; 0.10 mol) is added to the solution, and the resulting solution is stirred 18 hours at reflux under a nitrogen atmosphere. Cooling the red/brown solution to -5 °C in an ice-acetone bath causes the formation of a precipitate which is collected by filtration and washed with a small portion of cold nitromethane. The resulting tan solid is recrystallized from hot ethylene dichloride yielding the product as white crystals (1.8 g; 56% theory), m.p. 238-241 °C (dec.).

By utilizing the appropriate arylglycine in the procedure of Example 55 and following the procedure of this Example, the following 2-aryl-5-(trifluoromethyl)pyrrole-3-carbonitrile were prepared:



R	L	m.p. °C
H	H	215-218
H	4-CH ₃	191-193
H	4-OCH ₃	168-180(dec.)
3-Cl	4-Cl	245-246(dec.)
H	4-CF ₃	218-219

EXAMPLE 58

Preparation of 4-Bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile


Under a nitrogen purge, a suspension of 2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile (1.6 g; 0.005 mol) in acetic acid (25 mL) is heated, all the material dissolving to a clear solution at about 60 °C. A solution of bromine (0.8 mL; 0.015 mol) in acetic acid (10 mL) is added dropwise over 15 minutes to the refluxing solution. The solution is refluxed 6 hours then allowed to stir 18 hours at room temperature. The HPLC of the reaction mixture shows about 80% conversion to product. The mixture is heated back to reflux and more bromine (0.5 mL; 0.01 mol) in acetic acid (5 mL) is added dropwise. After refluxing another 3

hours, the aliquot shows > 95% conversion to product. The reaction is cooled, and solvent removed under reduced pressure on a rotary evaporator to obtain a dark grey solid. Toluene is added to the mixture and removed under reduced pressure, but the odor of acetic acid still remains. The entire material is dissolved in hot toluene (75 mL) to a turbid solution which is treated with DARCO filter and filtered. The light pink solution deposits a white solid upon cooling to ambient. After cooling in the freezer, the solid is collected by filtration, washed with hexanes, and dried on the filter. Further drying in a vacuum oven at 45 °C provides the product (1.2 g; app. 60% theoretical); m.p. 247-250 °C(dec.).

Anal. Calcd for $C_{12}H_5BrClF_3N_2$:

	C, 41.20;	H, 1.43;	N, 8.01;	Br, 22.89;	Cl, 10.16;	F, 16.31.
Found:	C, 41.27;	H, 1.48;	N, 8.10;	Br, 22.92;	Cl, 10.16;	F, 16.03.


By brominating the appropriate 2-aryl-5-(trifluoromethyl)pyrrole-3-carbonitrile, obtained by the procedure of Example 57, according to the above recipe, the following additional examples are prepared:



R	L	m.p. °C
H	H	235-238
H	4-CH ₃	244-245
3-Cl	4-Cl	218-223
H	4-CF ₃	225-226

EXAMPLE 59


Preparation of 2-(4-chlorophenyl)-5-trifluoromethyl-pyrrole-3,4-dicarbonitrile



Trifluoroacetic anhydride (3.1 mL; 0.022 mol) is added in a single portion to (4-chlorophenyl)glycine (2.0 g; 0.011 mol), causing an immediate yellow color and some refluxing. The mixture is slowly heated to reflux, causing all the material to dissolve to a yellow/orange solution which is heated 2 hours further. The reaction mixture is cooled, and solvent removed under reduced pressure. Toluene, is twice added and removed under reduced pressure to yield a very thick oil ($V_{CO} = 1800\text{cm}^{-1}$). This residue is dissolved (some insolubles) in CH_3NO_2 (40 mL) and bromofumaronitrile (2.7 g; 0.018 mol) is added in a single portion. The resulting solution is heated at reflux 18 hours, yielding a dark red solution. Solvent is removed under reduced pressure and the dark residue is dissolved in CH_2Cl_2 , some insolubles being removed by filtration. The material is fractionated via dry column chromatography (silica gel; 3% 2-PrOH in CH_2Cl_2), and appropriate fractions are taken. Evaporation of one fraction yields the desired compound as a yellow solid which is recrystallised from CH_3CN (DARCO treatment) to yield a pale yellow solid (0.2 g). m.p. = 238-241 °C. (some dec).

EXAMPLE 60

p-chloro- β -[(formylmethyl)aminol]cinnamitrile, diethyl acetal



A magnetically stirred solution of 250.0g (1.39 mol.) of p-chlorobenzoylacetonitrile, 203 mL (185.9g, 1.39mol) of 2,2-diethoxyethylamine, and 1300 mL of dried toluene is heated at reflux for 20 hours. Water is


collected in a Dean-Stark trap (23.8 mL, 95.2% theory). The hot cloudy dark brown solution with a large amount of undissolved solids is filtered through diatomaceous filter aid. After dilution with 200 mL of EtOAc, the solution is filtered through a 7cm X 13.5cm column of silica gel. The filtrate is concentrated in vacuo to give 354.3g (86.4% crude yield) of a clear dark oil which slowly solidifies. This solid is recrystallized from hot cyclohexane to give 324.2g (79.1% yield) of a waxy orange solid. NMR of this product shows it to be composed of 78% (Z) and 23% (E) isomeric mixture of p-chloro-β-[(formylmethyl)amino]cinnamionitrile, diethyl acetal, m.p. 60-72 °C. The following analytical data is for another similarly prepared sample.


Max(mull,Nujol): 3325(s), 3065(m), 2197(s), 1600(s), 1530(s), 1314(m), 1265(m), 1173(m), 1154(m), 1128(s), 1100(s), 1060(s), 1022(s), 939(m), 895(m), 844(s), 768(m), 730(m) Cm^{-1} .

¹H-NMR(chloroform): δ7.47 (d, J=8.6Hz, 2.12H, two aromatic protons), δ7.37 (d, J=8.6Hz, 2.12H, two aromatic protons), δ5.10(E) & δ4.86(Z) [br t, 1.25H, one N-H proton], δ4.69(Z) & δ4.60(E) [t, J=5.1Hz, 1.05H, one methine proton at the acetal carbon], δ4.07 (E) & δ4.05(Z) [s, 0.83H, enamine β proton], δ3.71(E) & δ3.68(Z) [q, J=7.1Hz, 2.22H, two methylene protons of one of two ethoxy groups], δ3.56(Z) & δ3.53(E) [q, J=7.1Hz, 2.22H, two methylene protons of one of two ethoxy groups], δ3.18 (t, J=5.1Hz, 1.77H, two methylene protons of the ethyleneacetal group), δ1.20 (t, J=7.1Hz, 4.90H, six methyl protons of the two ethoxy groups). ¹³C-NMR(chloroform): δ161.21 (α-enamine carbon), δ136.29 (Z) & δ134.60(E) [either C-1 or C-4 of the phenyl ring], δ134.08(Z) & δ132.30(E) [either C-1 or C-4 of the phenyl ring], δ129.34(Z) & δ129.89(E) [either C-2,6 or C-3,5 of the phenyl ring], δ128.94(Z) & δ128.63(E) [either C-2,6 or C-3,5 of the phenyl ring], δ121.19(Z) & δ119.50(E) [nitrile carbon], δ99.43(Z) & δ100.63(E) [β-enamine carbon], δ61.88(Z) & δ63.25(E) [methine carbon of the acetal], δ62.64(Z) & δ63.03(E) [methylene carbons of the ethoxy groups], δ46.32(Z) & δ47.33(E) [methylene carbon of the ethyl amine group], δ15.26 (methyl carbons of the ethoxy groups).


Microanalysis (Mw 294.78):				
Calcd:	C, 61.11%;	H, 6.50%;	N, 9.51%'	Cl, 12.03%.
Found:	C, 61.25%;	H, 6.25%;	N, 9.34%;	Cl, 12.35%.

Following the above procedure but substituting the appropriate benzoylacetoylacetonitrile for p-chlorobenzoylacetoneitrile and/or the appropriate 2,2-di(C₁-C₄ alkoxy)ethylamine for 2,2-diethoxyethylamine yields the following compounds:



L	M	R	(C ₁ -C ₄ alkoxy) ₂	mp °C
H	H	p-CH ₃ OCO	(OC ₂ H ₅) ₂	68-73
H	p-CH ₃	H	(OC ₂ H ₅) ₂	59-69
H	m-OCH ₃	p-OCH ₃	(OC ₂ H ₅) ₂	Red orange semi solid
H			(OC ₂ H ₅) ₂	62-70
p-Cl	H	H	(OCH ₃) ₂	---
H	p-CH ₃	H	(OCH ₃) ₂	---
H	m-Cl	p-Cl	(OC ₂ H ₅) ₂	---
H	H	p-OCF ₃	(OC ₂ H ₅) ₂	---
H	H	p-CF ₃	(OC ₂ H ₅) ₂	---

EXAMPLE 61

2(p-chlorophenyl)-pyrrole-3-carbonitrile

EP 0 347 488 B1

To 108 mL of trifluoroacetic acid stirred at 23 °C is added 54.00g (0.183mol) of solid p-chloro-β-[(formylmethyl)amino]cinnamionitrile, diethyl acetal over a period of 45 minutes. This addition produced an exotherm to 38 °C and, 32 minutes into the addition, a solid started to precipitate. After stirring at room temperature for 30 minutes, the reaction mixture is vacuum filtered and the collected solid is washed first with trifluoroacetic acid, secondly with an ethyl acetate-hexane mixture, and finally with hexane. The yield is 16.83g (45.4%) of an off-white solid, mp 165-166 °C. The following anal. data is from a similarly prepared sample.


Max(mull, Nujol): 3275(br s), 2225(s), 1502(s), 1410(m), 1275(m), 1200(m), 1108(s), 1023(m), 999(m), 908(m), 843(s), 752(s), 722(s), 695(s), 620(s) Cm^{-1} .

¹H-NMR(acetone): δ11.22 (v br s, 0.99H, one pyrrole N-H proton), δ7.82 (d, J=8.9Hz, 2.46H, two aromatic phenyl protons), δ7.51 (d, J=8.9Hz, 2.46Hz, two aromatic phenyl protons), δ7.02 (t, J=2.6Hz, 1.01H, one pyrrole proton at C-5), δ6.58 (t, J=2.6Hz, 0.77H, one pyrrole proton at C-4).

¹³C-NMR(acetone): δ137.73 (pyrrole C-2), δ134.42 (p-chlorophenyl at C-4), δ129.93 (methine carbons at C-3,5 of the phenyl ring), δ128.07 (methine carbons at C-2,6 of the phenyl ring), δ121.21 (pyrrole at C-5), δ117.93 (nitrile carbon), δ113.78 (pyrrole carbon at C-4), δ90.86 (pyrrole carbon at C-3).

Microanalysis (MW 202.64):				
Calcd.:	C, 65.19%;	H, 3.48%;	N, 13.83%;	Cl, 17.50%
Found:	C, 64.18%;	H, 3.52%;	N, 13.63%;	Cl, 17.74%

Use of the above procedure as shown or with the substitution of concentrated hydrochloric acid for trifluoroacetic acid affords the following compounds:




M and/or R	mp °C	Acid Used
4-Cl	165-166	conc. HCl, CF ₃ COOH
3,4-di-Cl	216-221	CF ₃ COOH
2-Cl	156-157	CF ₃ COOH
4-OCF ₃	143-145	CF ₃ COOH
4-CF ₃	179-180	CF ₃ COOH
2,4-di-Cl	197-199	CF ₃ COOH
3-Cl	150-156	CF ₃ COOH
4-CN	210-212	CF ₃ COOH
4-F	167-170	conc. HCl
4-SO ₂ CH ₃	221-221.5	CF ₃ COOH
3,4-di-F	173-175.5	CF ₃ COOH
3-CF ₃	166-168	CF ₃ COOH
4-COOCH ₃	155.5-158	CF ₃ COOH
4-CH ₃	117-137	CF ₃ COOH
4-NO ₂	174-177	CF ₃ COOH

EXAMPLE 62

4,5-Dichloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile

5



15

To a mechanically stirred solution of 16.83g (83.1mmol) of 2-(p-chlorophenyl)pyrrole-3-carbonitrile in 450 mL of glacial acetic acid at 36°C is added dropwise 14.7 mL (24.70g, 183.0mmol) of sulfonyl chloride over a period of 18 minutes. The addition produces a slight exotherm to 39°C and, after another 16 minutes, the reaction mixture is vacuum filtered. The collected solids are washed first with acetic acid and then with water. This solid after recrystallization from hot ethyl acetate, melts at 259-261°C. By similar procedures other samples of this product were prepared and the analytical data for one such product is shown below.

25

Max(mull, Nujol): 3170(br s), 3100(m), 2225(s), 1508(m), 1097(m), 825(s), 717(m), 660(m) cm⁻¹.

¹H-NMR(DMSO): δ7.72 (d, J = 8.6Hz, 2.00H, two aromatic protons), δ7.56 (d, J = 8.6Hz, 2.00H, two aromatic protons).

¹³C-NMR(DMSO): δ136.01 (pyrrole C-2 carbon), δ133.92 (p-chlorophenyl C-4 carbon), δ129.09 (p-chlorophenyl C-3,5 carbons), δ127.41 (p-chlorophenyl C-4 carbon), δ127.11 (p-chlorophenyl C-1 carbon), δ114.49 (nitrile carbon), δ114.10 (pyrrole C-5 carbon), δ110.92 (pyrrole C-4 carbon), δ90.09 (pyrrole C-3 carbon).

30

Microanalysis (MW 271.54):				
Calcd.:	C, 48.65%;	H, 1.86%;	N, 10.32%;	Cl, 39.17%
Found:	C, 49.22%;	H, 2.12%;	N, 9.85%;	Cl, 39.03%


35

EXAMPLE 63

40

4,5-Dibromo-2-(α,α,α-trifluoro-p-tolyl)-pyrrole-3-carbonitrile

45



50

55

EP 0 347 488 B1

To a stirred mixture of 0.8g of 2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile in 70 mL of chloroform is added 2 mL of bromine. The mixture, on stirring overnight, deposits a white solid which is collected by filtration. Thin layer chromatography (1:1 ethyl acetate-hexane) shows a single component; m.p. $>230^\circ\text{C}$.


5

Anal. Calc'd for $\text{C}_{12}\text{H}_5\text{Br}_2\text{F}_3\text{N}_2$;				
	C, 36.55;	H, 1.27;	N, 7.11;	Br, 40.61.
Found:	C, 36.40;	H, 1.08;	N, 6.99;	Br, 40.55.

10

Following the above procedure but substituting the appropriately substituted phenylpyrrole-3-carbonitrile for 2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile yields the following compounds.

15



20

25

30

35

40


45

50

55

L	M	R	X	Y	mp °C
H	H	4-NO ₂	Br	Br	274-277
H	H	4-F	Cl	Cl	>220
H	H	4-F	Br	Br	>220
H	H	4-SO ₂ CH ₃	Cl	Cl	>230
H	3-F	4-F	Cl	Cl	>230
H	3-F	4-F	Br	Br	>220
2-Cl	3-Cl	4-Cl	Cl	Cl	
2-Br	3-Br	4-Br	Br	Br	
H	H	4-OCF ₃	Cl	Cl	222-225
H	H	4-OCF ₃	Br	Br	
H	H	4-OCF ₃	Cl	H	
H	H	4-CN	Br	Br	>230
H	H	4-CN	Cl	Cl	>240
H	H	4-SO ₂ CH ₃	Br	Br	>230
H	H	4-NO ₂	Cl	Cl	246-249
H	3-Cl	4-Cl	Br	Br	>260
H	H	3-CF ₃	Cl	Cl	>230
H	H	4-COCH ₃	Cl	Cl	251-254
H	2,3-CH=CH-		Cl	Cl	244-247
H	H	4-CH ₃	Cl	Cl	215-217
H	2-Cl	4-Cl	Br	Br	>230
H	H	3-Cl	Cl	Cl	>230
H	2-Cl	4-Cl	Cl	Cl	>230
H	H	4-Cl	Br	Br	273-274
H	H	2-Cl	Br	Br	>230
H	H	4-CF ₃	Cl	Cl	>230
H	H	4-Br	Cl	Cl	>235
H	H	2-Cl	Cl	Cl	>230
H	3-Cl	4-Cl	Cl	Cl	>235
H	H	H	Cl	Cl	254-255


EXAMPLE 64


 α -(2,2-diethoxyethylamino)- β -nitrostyrene and 3-nitro-2-phenylpyrrole


Alpha-nitro acetophenone (5.7g, 0.0345m) is taken up in 100 mL toluene and 4.6g (0.0345m) of amino acetaldehyde diethyl acetal is added. The reactants are put into a 250 mL RB flask fitted with a Dean-Stark trap. The trap is filled with 4A molecular sieves and the mixture is heated at reflux for 18 hours. The toluene is removed in vacuo to give 8.36g of α -(2,2-diethoxyethylamino)- β -nitrostyrene as a brown oil. To this oil is added 50 mL of concentrated HCl. As the flask is swirled the oil turns to a yellow suspension. After 10 minutes the solid is filtered to give 2.48g of a yellow solid. Recrystallization from ether/ethylacetate/hexane gives the product as two fractions, 2.08g of m.p. 190-192 °C, (31%).

Max 1485 cm⁻¹(NO₂), H-NMR(CDCl₃/DMSO) δ 6.73(m,2H), 7.46(m,5H).

Other β -nitrostyrene compounds can be prepared by the above reaction by substituting the appropriately substituted α -nitro acetophenone for α -nitro acetophenone and/or appropriate 2,2-di(C₁-C₄ alkoxy)-ethylamine for amino acetaldehyde diethyl acetal to give the following compounds



L	M	R	(C ₁ ---C ₄ alkoxy) ₂
H	H	p-CH ₃ OCO	(OC ₂ H ₅) ₂
H	p-CH ₃	H	(OC ₂ H ₅) ₂
H	m-OCH ₃	p-OCH ₃	(OC ₂ H ₅) ₂
H			(OC ₂ H ₅) ₂
p-Cl	H	H	(OCH ₃) ₂
H	p-CH ₃	H	(OCH ₃) ₂
H	H	p-CF ₃	(OC ₂ H ₅) ₂

EXAMPLE 652,3-Dichloro-4-nitro-5-phenylpyrrole

55 A mixture of 3-nitro-2-phenylpyrrole (1.56g, 0.0083m) in 60 mL of dioxane is cooled in an ice bath while 25.9g (.0182m) of commercial sodium hypochlorite is added dropwise. After stirring for 45 minutes, the mixture is acidified with concentrated HCl. Water and Et₂O are added. The layers are separated and the top organic layer is washed with H₂O, dried over anhydrous MgSO₄ and concentrated in vacuo to give 2.21g of yellow solid. Purification by chromatography using silica gel and eluting with increasing ratios of ethyl

acetate/hexane gives, after stripping, 0.77g of yellow solid (36%) m.p. 190-190.5 °C;

Analysis: Calcd. for C ₁₀ H ₆ N ₂ O ₂ Cl ₂			
Found:	C, 46.72;	H, 2.35;	N, 10.90
	C, 46.96;	H, 2.06;	N, 10.02

EXAMPLE 66

Insecticide and acaricide evaluations

All tests are preformed using technical materials. All concentrations reported herein are in terms of active ingredient. All tests are kept at 27 °C.

Spodoptera eridania, 3rd instar larvae, southern armyworm

A Sieva lima bean leaf expanded to 7-8 cm in length is dipped in the test suspension with agitation for 3 seconds and placed in a hood to dry. The leaf is then placed in a 100x10 mm petri dish containing a damp filter paper on the bottom and ten 3rd instar caterpillars. The dish is maintained for 5 days before observations are made of mortality, reduced feeding, or any interference with normal moulting.

Spodoptera eridania, 7-day residual

The plants treated in the above Test are maintained under high intensity lamps in the greenhouse for 7 days. These lamps duplicate the effects of a bright sunny day in June in New Jersey and are kept on for 14 hour day length. After 7 days, the foliage is sampled and assayed as in the above-said Test.

Aphis fabae, mixed instar, bean aphid

Pots containing single nasturtium plants (*Tropaeolum* sp.) about 5 cm tall are infested with about 100-200 aphids one day before the test. Each pot is sprayed with the test formulation for 2 revolutions of a 4 rpm turntable in a hood, using a #154 DeVilbiss atomizer. The spray tip is held about 15 cm from the plant and the spray directed so as to give complete coverage of the plants and the aphids. The sprayed pots are set on their sides on white enamel trays and held for 2 days, following which mortality estimates are made.

Tetranychus urticae (P-resistant strain), 2-spotted spider mite

Sieva lima bean plants with primary leaves expanded to 7-8 cm are selected and cut back to one plant per pot. A small piece is cut from a leaf taken from the main colony and placed on each leaf of the test plants. This is done about 2 hours before treatment to allow the mites to move over to the test plant and to lay eggs. The size of the cut piece is varied to obtain about 100 mites per leaf. At the time of the treatment, the piece of leaf used to transfer the mites is removed and discarded. The mite-infested plants are dipped in the test formulation for 3 seconds with agitation and set in the hood to dry. Plants are kept for 2 days before estimates of adult kill are made using the first leaf. The second leaf is kept on the plant for another 5 days before observations are made of the kill of eggs and/or newly emerged nymphs.

Diabrotica undecimpunctata howardi, 3rd instar southern corn rootworm

One cc of fine talc is placed in a 30 ml wide-mouth screw-top glass jar. One ml of the appropriate acetone suspension is pipetted onto the talc so as to provide 1.25 and 0.25 mg of active ingredient per jar. The jars are set under a gentle air flow until the acetone is evaporated. The dried talc is loosened, 1 cc of millet seed is added to serve as food for the insects and 25 ml of moist soil is added to each jar. The jar is capped and the contents thoroughly mixed on a Vortex Mixer. Following this, ten 3rd instar rootworms are added to each jar and the jars are loosely capped to allow air exchange for the larvae. The treatments are held for 6 days before mortality counts are made. Missing larvae are presumed dead, since they decompose rapidly and can not be found. The concentrations used in this test correspond approximately to 50 and 10 kg/ha, respectively.

Rating Scale:	
0 = no effect	5 = 56-65% kill
1 = 10-25% kill	6 = 66-75% kill
2 = 26-35% kill	7 = 76-85% kill
3 = 36-45% kill	8 = 86-99% kill
4 = 46-55% kill	9 = 100% kill
	R - reduced feeding

Table I

Compound	BEAN	ARHWORMS			P. RES	MITES	SCRW
	APHIDS	ppm	1000	ppm	ppm	ppm	kg/ha
4,5-dichloro-2-phenylpyrrole-3-carbonitrile	100	0	9	100	300	50	0
4,5-dichloro-2-(p-chlorophenyl)-pyrrole-3-carbonitrile	0	0	9	8.5	0	5.5	0

Table I (Cont.)

	BEAN APHIDS	ARMYWORMS			P. RES MITES	SCRW
	ppm	ppm	ppm	days	ppm	kg/ha
Compound	100	1000	100	7	300	50
4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	9	9	0	0
4,5-dichloro-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	7.5	9	9	9	9	7.7

Table I (Cont.)

Compound	BEAN	ARMYWORMS			P. RES	SCRW
	APHIDS	ppm	ppm	7 days	MITES	kg/ha
4,5-dichloro-2-(o-chlorophenyl)-pyrrole-3-carbonitrile	100	1000	100		300	50
	0	9	9		0	0
2-(p-bromo-phenyl)4,5-dichloropyrrole-3-carbonitrile	0	9	9		8	0
4,5-dichloro-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	0	9	9		9	8

Table I (Cont.)

	BEAN APHIDS	ARMYWORMS		P.RES MITES	SCRW
	ppm	ppm	7 days	ppm	kg/ha
Compound	100	1000	100	300	50
4,5-dibromo-2-(o-chlorophenyl)- pyrrole-3-carbo- nitrile	0	9	4	0	0
4,5-dibromo-2-(p-chlorophenyl)- pyrrole-3-carbo- nitrile	0	9	9	0	0

Table I (Cont.)

Compound	BEAN	ARMYWORMS			P.RES	SCRW
	APHIDS				MITES	
	ppm		ppm	days	ppm	kg/ha
4,5-dibromo-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	100	1000	100	7	300	50
	0	9	9		9	9
4,5-dichloro-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	9	9	0	0

Table I (Cont..)

Compound	BEAN	ARMYWORMS			P. RES	SCRW
	APHIDS				MITES	
	ppm		ppm	days	ppm	kg/ha
4,5-dibromo-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile	100	1000	100	7	300	50
	0	9	9	9	0	0
4,5-dichloro-2-(m-chlorophenyl)-pyrrole-3-carbonitrile	0	9	9	9	0	0

Table I (Cont.)

Compound	BEAN	ARMYWORMS			P.RES	SCRW
	APHIDS	ppm	ppm	days	MITES	kg/ha
2,3-dichloro-4-nitro-5-phenylpyrrole	100	1000	100	7	300	50
	0	9	9	9	0	0
2,3-dichloro-5-(p-chlorophenyl)-4-nitropyrrole	0	9	9	9	7.5	8
2,3-dibromo-5-(p-chlorophenyl)-4-nitropyrrole	0	9	9	9	0	7.5

Table I (Cont.)

	BEAN APHIDS	ARMYWORMS			P. RES MITES	SCRW
	ppm	1000	ppm	7 days	ppm	kg/ha
Compound	100		100		300	50
2,3-dibromo-4-nitro-5-phenylpyrrole	0	9	9	9	0	0
2,3-dichloro-5-(3,4-dichloro-phenyl)-4-nitropyrrole	0	9	9	9	0	0

Table I (Cont.)

Compound	BEAN APHIDS		ARMYWOORMS		P. RES MITES	SCRW
	ppm		ppm	days	ppm	kg/ha
2-(p-bromo-phenyl) 4,5-dichloro-3-nitropyrrole	100	1000	100	7	300	50
	8	9	9	9	8.5	8
2,3-dichloro-4-nitro-5-(alpha,alpha-trifluoro-p-tolyl)pyrrole	8.3	9	9	9	9	8.3

EXAMPLE 67Insecticidal evaluationsHeliothis virescens, 3rd instar tobacco budworm

Cotton cotyledons are dipped in the test formulation and allowed to dry in a hood. When dry, each is cut into quarters and ten sections placed individually in 30 ml plastic medicine cups containing a 5-7 mm long piece of damp dental wick. One third-instar caterpillar is added to each cup and a cardboard lid placed

on the cup. Treatments are maintained for 3 days at before mortality counts and estimates of reduction in feeding damage are made.

Empoasca abrupta, adults, western potato leafhopper

5

A Sieva lima bean leaf about 5 cm long is dipped in the test formulation for 3 seconds with agitation and placed in a hood to dry. The leaf is placed in a 100x10 mm petri dish containing a moist filter paper on the bottom. About 10 adult leafhoppers are added to each dish and the treatments are kept for 3 days before mortality counts are made.

10

Blattella germanica, bait test, adult male German cockroach

A 0.1% bait is prepared by pipetting 1 ml of a 1000 ppm solution of the test compound in acetone onto 1 gram of cornmeal in a 30 ml wide-mouth bottle. The bait is dried by passing a gentle stream of air into the bottle. The bait is placed in a 1 pint wide-mouth Mason jar and ten adult male cockroaches are added. A screen lid is placed on the jar and a small piece of cotton soaked in 10% honey is put on the top of the screen lid. Mortality counts are made after 3 days.

20

Blattella germanica, residue test, adult male German cockroach

One ml of a 1000 ppm acetone solution of the test material is pipetted slowly over the bottom of a 150 x 15 mm petri dish so as to give as uniform coverage as possible. After the deposit has dried, 10 adult male cockroaches are placed in each dish and the lid is added. Mortality counts are made after 3 days.

25

Scodoptera eridania, systemic uptake, 3rd instar larvae, southern armyworm

The compound is formulated as an emulsion containing 0.1 gm of the test material, 0.2 gm of Emulphor EL-620® emulsifier, 10 ml of acetone and 90 ml of water. This is diluted 10-fold with water to give a 100 ppm emulsion for the test. Subsequent 10-fold dilutions are made with water as needed. Sieva lima bean plants, with the primary leaves expanded to a length of 7-8 cm, are cut off at least 3 cm above the soil level to avoid contamination with soil bacteria that will cause decay of the stem during the test. The cut stems are placed in the test emulsions and each stem is wrapped with a bit of cotton to hold the stem off the bottom of the bottle and to limit evaporation and volatilization of the compound. The test is maintained for 3 days at 27°C to allow the compounds to be taken up into the plant. Following this, one leaf is removed from the plant and placed in a 100 x 10 mm petri dish with 10 southern armyworms as described in Test III. Mortality counts and observations of feeding damage are made 3 and 5 days later.

Empoasca abrupta, Adults, Western Potato Leafhoppers, Systemic Uptake

40

The compound is formulated as an emulsion containing 0.1 gm of the test material, 0.2 gm of Emulphor ES-620® emulsifier, 10 ml of acetone and 90 ml of water. This is diluted 10-fold with water to give a 100 ppm emulsion for the test. Subsequent 10-fold dilutions are made with water as needed. Sieva lima bean plants, with the primary leaves expanded to a length of 7-8 cm, are cut off at least 3 cm above the soil level to avoid contamination with soil bacteria that will cause decay of the stem during the test. The cut stems are placed in the test emulsions and each stem is wrapped with a bit of cotton to hold the stem off the bottom of the bottle and to limit evaporation and volatilization of the compound. The test is maintained for 3 days at 27°C to allow the compounds to be taken up into the plant. Following this, one leaf is removed from the plant and placed in a 100 x 10 mm petri dish and tested as in Test VIII, above.

50

55

The rating scale for the above tests is the same as described in Example 9.

Table II

	<u>LEAF</u>		<u>TBW</u> ³		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>				<u>LEAF</u>		<u>RES.</u>	
	ppm	100	1000	100	100	100	1000	1000
					<u>SAW</u>	<u>HOPPER</u>	<u>BAIT</u>	
					ppm	ppm	ppm	ppm
4,5,-dichloro-2-phenylpyrrole-3-carbonitrile	0	8		0	-	0	0	0
4,5-dichloro-2-(p-chlorophenyl)-pyrrole-3-carbonitrile	0	9	8.5	9	0	0	0	7

Table II (Cont.)

	<u>LEAF</u>	<u>TBW</u> ³	<u>C-S SYSTEMIC</u>	<u>G. COCKROACH</u>
	<u>HOPPER</u>			
			<u>LEAF</u>	<u>RES.</u>
			<u>SAW</u>	<u>BAIT</u>
			<u>HOPPER</u>	
	ppm	ppm	ppm	ppm
	100	1000	100	1000
4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	9	7
4,5-dichloro-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	9	9	9	8

Table II (Cont.)

	<u>LEAF</u>		<u>TBW</u> ³		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>				<u>SAW</u>	<u>LEAF HOPPER</u>	<u>BAIT</u>	<u>RES.</u>
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm
	100	1000	100	100	100	100	1000	1000
4,5-dichloro-2-(o-chlorophenyl)-pyrrole-3-carbonitrile	0		0		7	0	0	0
2-(p-bromophenyl)-4,5-dichloropyrrole-3-carbonitrile	0	9	9		-	-	0	0

Table II (Cont.)

	<u>LEAF</u>		<u>TBW</u> ³	<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>		
	<u>HOPPER</u>			<u>SAW</u>		<u>LEAF HOPPER</u>		
	ppm	100	1000	ppm	100	1000	RES. ppm	1000
4,5-dichloro-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	9	-	9	0	9	
4,5-dibromo-2-(o-chlorophenyl)-pyrrole-3-carbonitrile	0		0	7	0	0	0	

Table II (Cont.)

	<u>LEAF</u>		<u>TBW³</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>							
	ppm	1000	ppm	100	ppm	100	ppm	1000
	0	9	9	9	-	0	0	0
4,5-dibromo-2-(p-chlorophenyl)-pyrrole-3-carbonitrile								
4,5-dibromo-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	9	9	-	9	0	9

Table II (Cont.)

	<u>LEAF</u>		<u>TBW</u> ³	<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>			<u>SAW</u>	<u>LEAF HOPPER</u>	<u>BAIT</u>	<u>RES.</u>
	ppm	ppm	ppm	ppm	ppm	ppm	ppm
	100	1000	100	100	100	1000	1000
4,5-dichloro-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	9	-	0	0	0
4,5-dibromo-2-(2,4-dichlorophenyl)pyrrole-3-carbonitrile	0		0	-	0	0	0

Table II (Cont..)

	<u>LEAF</u>		<u>TBW³</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>				<u>SAW</u>	<u>HOPPER</u>	<u>BAIT</u>	<u>RES.</u>
	ppm	100	1000	ppm	100	100	1000	ppm
	100	1000	100	100	100	100	1000	1000
4,5-dichloro-2-(m-chlorophenyl)-pyrrole-3-carbonitrile	0	9	9	-	0	0	0	0
2,3-dichloro-4-nitro-5-phenylpyrrole	0	9	6	8	9	0	0	4

Table II (Cont.)

	<u>LEAF</u>		<u>TBW³</u>	<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>			<u>SAW</u>	<u>LEAF HOPPER</u>	<u>BAIT</u>	<u>RES.</u>
	ppm	ppm	ppm	ppm	ppm	ppm	ppm
	100	1000	100	100	100	1000	1000
2,3-dichloro-5-(p-chlorophenyl)-4-nitropyrrole	8.5	9	8	9	0	9	9
2,3-dibromo-5-(p-chlorophenyl)-4-nitropyrrole	0	8.5	6	0	0	0	9
2,3-dibromo-4-nitro-5-phenylpyrrole	0	8.5	0	9	0	0	0

Table II (Cont.)

	<u>LEAF</u>		<u>TBW³</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>				<u>SAW</u>	<u>LEAF HOPPER</u>	<u>BAIT</u>	<u>RES.</u>
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm
	100	1000	100	100	100	100	1000	1000
2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole	9	9	9	9	9	9	0	9
2-(p-bromo-phenyl)-4,5-dichloro-3-nitropyrrole	8	9	6.5	9	9	9	0	9

Table II (Cont.)

	<u>LEAF</u>		<u>TBW³</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>				<u>SAW</u>	<u>LEAF</u> <u>HOPPER</u>	<u>BATT</u>	<u>RES.</u>
	ppm		ppm		ppm	ppm	ppm	ppm
	100	1000	100	100	100	100	1000	1000
	9	9	8	9	9	9	0	9
2,3-dichloro-4-nitro-5-alpha,alpha-trifluoro-p-tolyl)pyrrole								

EXAMPLE 68

A) Evaluation of test compounds as nematicidal agents Culture Maintenance: Cultures of *C. elegans* (Bristol strain from J. Lewis) are maintained on *E. coli* lawns on NG Agar Plates at 20°C. New cultures are established weekly.

Nematodes for testing are washed from 4-5 day old cultures using Fresh Ascaris Ringers Solution (FARS). The worms are further washed with FARS, containing gentamycin, to reduce bacterial contamination and centrifuged to separate worms from wash solution. This procedure is repeated three times. The washed worms are then added to *C. briggsae* Maintenance Medium (CbMM), from GIBCOa to which is added gentamycin (600 units/ml) and mycostatin (0.5 mg/ml).

The tests are then made with mixtures of three compounds, piggy-backed from another high capacity screening program to reduce additional labor and compound expenditures.

Compounds are dissolved in acetone and made up to volume with equal parts of water. The final test concentration of each compound in the mixture is 150 ppm. The test material is micropipetted (25 ul) into a single well of a 96-well sterile tissue culture plate (COSTAR)^b and the solvent allowed to evaporate. These "treated" plates are used immediately or stored in a freezer without apparent adverse effects on the compounds.

A freshly prepared volume (50 ug) of *C. elegans* in CbMM is micropipetted into each treated well and several control wells per plate. Culture plate are incubated at 20 ° C.

Observations for efficacy are made under a dissecting microscope at 4, 24 and 48 hours post-immersion. Immediately prior to reading the plate, it is gently tapped to stimulate the movement of the worms. Activity is judged subjectively, but semi-quantitatively, based on the drug effects on motility of the adults and larvae. The criteria are as follows: 8 = no motility, 7 = markedly reduced motility in approximately 95% of worms, 6 = reduced motility, 5 = slightly reduced motility, 0 = normal motility, same as controls. Other factors indicating activity are easily noted such as death, rigor mortis, contraction, coiling, paralysis, abnormal twitching, reduced worm population in 48 hours and other deviation from normal behavior.

	PROCEDURE FOR <u>CAENORHABDITIS ELEGANS</u> ASSAY	
20	Day 0	.Inoculate <u>E. Coli</u> -NG Agar Dish With 30-50 <u>C. Elegans</u> .Incubate At 20 ° C.
25	Day 4	.Harvest New <u>C. Elegans</u> Population .Wash With Antibiotics .Transfer To CbMM .Add <u>C. Elegans</u> (25-100 UL) To "Medicated" Wells ^a .Observe For Activity At 4 Hours Post-Immersion
30	Day 5 Day 6	.Observe For Activity .Observe For Activity

^a Medicated Wells May Be Prepared Fresh Or Earlier And Stores In Freezer

Data obtained in these tests are reported in Table III below.

35 B) Root-Knot Nematode Assay

Populations of the root-knot nematode (Meloidogyne incognita) are maintained on Fireball tomatoes in the greenhouse. Egg masses are removed from the infested root surfaces and are kept on moistened filter paper for 24 hours to allow them to hatch. Larvae emerge and drop into the water beneath the paper. Larvae for test are transferred to cell plate wells containing test compounds at 300 ppm in 3% acetone, about 10 larvae per cell well. Infested wells are held at 27 ° C and mortality is determined 24 hours after treatment.

Data obtained are reported in Table III below.

Table III

5		C. Ele. 150 ppm		Root Knot Nematodes. 300 ppm
		L	A	
10	4,5-dichloro-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	-	-	4
	4,5-dichloro-2-(alpha, alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	5
	4,5-dibromo-2-(alpha, alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	0
	2,3-dichloro-4-nitro-5-phenylpyrrole	0	0	9
	2,3-dichloro-5-(p-chlorophenyl)-4-nitropyrrole	9	9	9
15	2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole	9	9	0
	2-(p-bromophenyl)-4,5-dichloro-3-nitropyrrole	9	9	6
	2,3-dichloro-4-nitro-5-(alpha,alpha,alphatrifluoro-p-tolyl)pyrrole	9	9	-

EXAMPLE 69

Following the procedures of Examples 59 and 60, compounds of the invention are evaluated against a variety of insect species including: leaf hoppers, tobacco budworm, southern armyworm, and the German cockroach. The rating system is the same system used in the above-said examples. Data obtained are reported in Table IV below. Where two or more tests have been conducted with the same test compound, the results are overaged. also, a - in the table indicates no test.

Table IV

Compound	<u>LEAF</u> <u>HOPPER</u>		<u>TBW</u> ³		<u>ARMYFORMS</u>				<u>C-S SYSTEMIC</u>			<u>G. COCKROACH</u>		
	ppm	100	1000	ppm	100	1000	ppm	7	days	100	100	<u>LEAF</u> <u>HOPPER</u>		<u>RES.</u> ppm
												<u>SAW</u> ppm	<u>HOPPER</u> ppm	
2,5-dichloro-4-phenylpyrrole-3-carbonitrile	0	6	0	9	9	9	0	8	0	0	0	0	0	
2,3-dibromo-4-nitro-5-phenylpyrrole	0	9	0	9	9	9	9	-	0	9	0	0	0	
4-chloro-2-(p-chlorophenyl)-pyrrole-3-carbonitrile	0	9	2	9	9	9	9	-	9	9	9	0	0	

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	ppm	1000	ppm	100	ppm	1000	100	ppm	100	100	ppm	RES.
4,5-dichloro-2-(o-chlorophenyl)pyrrole-3-carbonitrile	0	0	0	0	9	9	9	7	7	0	0	0
5-bromo-2-(p-chlorophenyl)pyrrole-3-carbonitrile	0	9	0	0	9	9	9	-	-	0	0	0
4,5-dichloro-2-(3,4-dichlorophenyl)-1-(ethoxymethyl)pyrrole-3-carbonitrile	9	9	9	9	9	9	9	5	5	0	0	7

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	1000	ppm	100	ppm	1000	ppm	100	ppm	1000
4,5-dichloro-2-(p-chlorophenyl)-1-methylpyrrole-3-carbonitrile	0	9	9	9	9	9	-	0	9	9
p-(4,5-dichloro-3-cyanopyrrole-2-yl)methylbenzoate	0	4R8	0	0	9	9	3	0	0	0
4,5-dibromo-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	0	0	9	9	0	0	0	0

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS				C-S SYSTEMIC				G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYWORMS				C-S SYSTEMIC				G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	7 days	ppm	100	100	100	RES.	ppm
4,5-dichloro-2-(alpha, alpha-trifluoro-m-tolyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9	0	0	9	9
4,5-dichloro-2-(3,4-dichlorophenyl)-1-ethylpyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9	9	9	0	0
4,5-dichloro-2-(3,4-difluorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9	9	9	0	0

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS		C-S SYSTEMIC		G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYWORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	100	ppm	RES.
4,5-dichloro-2-(3,4-dichlorophenyl)-1-methylpyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9
4,5-dichloro-2-[p-(methylsulfonyl)phenyl]pyrrole-3-carbonitrile	0	-	0	7	0	0	9	9	9	0
4,5-dibromo-1-methyl-2-(alpha, alpha, alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	0	9	8	9	9	9	9	9	9	8.5

Table IV (Cont.)

Compound	<u>LEAF</u> <u>HOPPER</u>		<u>TBW</u> ³		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES. ppm
4,5-dichloro-2-(p-fluorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	0
4,5-dibromo-2-(3,4-difluorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	4.5	4.5	4.5	0
4,5-dibromo-2-(p-fluorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	0	0	0	0

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	7	SAW	LEAF HOPPER	RES.	ppm
		100	1000		100	1000		days	ppm	ppm		1000
4,5-dibromo-2-(p-nitrophenyl)pyrrole-3-carbonitrile	0	9	9	6	9	9	9	4	4.5	9		9
4,5-dichloro-2-(p-nitrophenyl)pyrrole-3-carbonitrile	0	9	9	7	9	9	9	9	4.5	4.5		9
1-benzyl-4,5-dibromo-2-(alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	9	9	9	9	9	9	9	6		0

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			LEAF HOPPER		G. COCKROACH	
	ppm		ppm		ppm			ppm			ppm		ppm	
	100	1000	100	1000	100	1000	1000	100	100	100	100	100	100	1000
4,5-dichloro-2-(p- cyanophenyl)pyrrole -3-carbonitrile	0	9	8	9	9	9	-	0	0	0	0	0	5	
4,5-dibromo-2-[p- (methylsulfonyl) phenyl]pyrrole-3- carbonitrile	0	2	-	9	0	9	-	0	0	0	0	0	0	
4,5-dibromo-2-(p- chlorophenyl)pyr- role-3-carbonitrile	0	9	7	9	9	9	-	0	0	0	0	0	0	

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	1000	ppm	100	ppm	1000	ppm	100	ppm	1000
4,5-dichloro-2-(3,4-dichlorophenyl)-1-[2-(methylthio)ethyl]pyrrole-3-carbonitrile	0	8		5	9	0	9	0	0	0
1-methyl-4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	7	7		0	9	0	9	0	0	0

Table IV (Cont.)

Compound	LEAF HOPPER		TPW ³		ARMYBORMS		C-S SYSTEMIC				G. COCKROACH	
							SAW		HOPPER		RES.	
	ppm	100	1000	ppm	100	1000	ppm	100	ppm	100	ppm	1000
4,5-dichloro-1-methyl-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	9	9	9	9	9	9	9	9	9	9
5-bromo-4-chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile	2.5	-	9	9	9	9	9	0	6	9	9	9
2,3-dichloro-5-(3,4-dichlorophenyl)-1-(ethoxymethyl)-4-nitropyrrole	6	5	0	9	9	9	9	9	9	9	9	9

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	1000	ppm	100	ppm	1000	7 days	ppm	100	1000
4-bromo-5-chloro-2-(p-chlorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	7	0	9
1-benzyl-4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	0	9	9	9	5.5	9	9	2
Ethyl 2,3-dichloro-5-(3,4-dichlorophenyl)-4-cyanopyrrole-1-acetate	0	9	4	9	9	9	7	9	0	0

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS		C-S SYSTEMIC		G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYWORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES. ppm
4,5-dichloro-1-(ethoxymethyl)-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	9	9	9	9	9	9	9	9	9	9
3-bromo-5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonyl nitrile	0	9	8.5	9	9	9	5	0	9	9
4,5-dichloro-2-(3,4-dichlorophenyl)-1-(2-propynyl)pyrrole-3-carbonitrile	8	9	9	9	9	9	9	9	9	9

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	7	SAW	LEAF	RES.	
								days	ppm	HOPPER	ppm	
									100	100	100	1000
4,5-dibromo-3-(p-chlorophenyl)pyrrole-2-carbonitrile	0	9	3	9	9	9	9	9	0	0	0	0
5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonitrile	7	9	9	9	9	9	9	9	5	9	7	7
5-bromo-4-chloro-(3,4-dichlorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	0	6	9	9

Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYWORMS			C-S SYSTEMIC			G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	7 days	100	100	100	RES. ppm
4,5-dibromo-3-(p-chlorophenyl)-1-methylpyrrole-2-carbonitrile	0	-	-	9	9	9	9	0	0	0	0	9
2-bromo-5-phenylpyrrole-3,4-dicarbonitrile	0	2	0	9	0	9	9	9	0	0	0	0
4-bromo-2-phenyl-5-(trifluoromethyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9	9	0

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYFORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES. ppm
							7	100	100	1000
							days	100	100	1000
2-(3,4-dichloro-phenyl)-5-nitro-pyrrole-3-carbonitrile	7	9	0	9	9	9	9	0	0	0
4-bromo-2-(3,4-dichlorophenyl)-5-nitropyrrole-3-carbonitrile	0	5	0	9	0	9	3	0	0	0
2,4-dibromo-5-phenylpyrrole-3-carbonitrile	0	9	9	9	9	9	9	0	0	0

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES. ppm
2-(3,4-dichloro-phenyl)-4,5-diiodo-pyrrole-3-carbo-nitrile	0	9	9	9	9	9	7	100	100	ppm
2,3-dibromo-5-(p-chlorophenyl)-1-(ethoxymethyl)-4-nitropyrrole	9	9	9	9	9	9	9	100	100	ppm
1-benzyl-2,3-dibromo-5-(p-chlorophenyl)-4-nitropyrrole	0	4	0	0	8	9	0	9	9	ppm

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	1000	ppm	100	ppm	1000	7	ppm	100	1000
2,3-dibromo-5-(p-chlorophenyl)-1-methyl-4-nitro-pyrrole	0	8		7	9	9	7	0	4	0
4,5-dibromo-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	9	9		9	9	9	9	9	0	0
4,5-dichloro-1-(ethoxymethyl)-2-[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	9	9		9	9	9	9	9	9	9

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARTEMIS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	1000	ppm	100	ppm	1000	SAW	LEAF HOPPER	RES.	
	100	1000	ppm	100	ppm	1000	ppm	ppm	ppm	1000
							7	days	100	100
5-(p-chloro-phenyl)-pyrrole-2,4-dicarbonitrile	3	9	9	9	9	9	9	9	0	8
1-benzyl-4,5-dichloro-2[p-(trifluoromethoxy)phenyl]pyrrole-3-carbonitrile	9	9	9	0	9	9	9	9	9	0
4,5-dichloro-2-(p-chlorophenyl)-1-(ethoxymethyl)pyrrole-3-carbonitrile	8.5	9	9	9	9	9	9	9	9	0

Table IV (Cont.)

Compound	LEAF HOPPER		TBA ³		ARITHORUS			C-S SYSTEMIC			G. COCKROACH	
	LEAF HOPPER		TBA ³		ARITHORUS			C-S SYSTEMIC			G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	7	SAW	LEAF HOPPER	RES.	ppm
	100	1000	100	1000	100	1000	100	days	100	100	100	1000
4,5-dichloro-2-(3,4-dichlorophenyl)-1-(2-hydroxyethyl)pyrrole-3-carbonitrile	0	9	2	9	9	9	0	9	0	9		0
2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile	7	9	0	9	0	9	0	7	0	9		0
4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)-pyrrole-3-carbonitrile	9	9	9	9	9	9	9	9	9	9		9

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	1000	ppm	100	ppm	1000	ppm	100	ppm	1000
	100	1000	ppm	100	ppm	1000	7 days	100	100	1000
4-bromo-2-(p-chlorophenyl)-5-nitropyrrole-3-carbonitrile	0	9	0	0	9	9	9	0	7	0
3-bromo-5-(p-chlorophenyl)pyrrol-2,4-dicarbonitrile	6	8	5	5	9	9	9	0	3	9
4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-1,3-dicarbonitrile	0	9	0	0	9	9	9	0	0	9

Table IV (Cont.)

Compound	LEAF HOPPER		TBA ³		ARMYAORMS		C-S SYSTEMIC		G. COCKROACH	
	LEAF HOPPER		TBA ³		ARMYAORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	100	ppm	100	ppm	1000	7	ppm	ppm	RES.
	100	1000	ppm	100	ppm	1000	days	100	100	1000
1-[(benzyloxy) methyl]-4,5- dichloro-2-(3,4- dichlorophenyl) pyrrole-3-carbo- nitrile	5	9		0		9	9	9	9	0
4,5-dichloro-2-(2, 4-dichloro-5-fluoro- phenyl)pyrrole-3- carbonitrile	4	5		0		9	9	9	0	9
5-[p-(trifluoro- methoxy)phenyl]pyr- role-2,4-dicarbo- nitrile	7	7		0		9	9	8	0	8

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW</u> ³	<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>								
	ppm	100	1000	ppm	100	1000	ppm	100	1000
4,5-dichloro-1- [(p-chlorophenoxy) methyl-2-(3,4- dichlorophenyl) pyrrole-3-carbo- nitrile	0	8	0	0	9	9	9	9	0
4,5-dichloro-2- (3,4-dichloro- phenyl)-1-(3-iodo- -2-propynyl)pyrrole -3-carbonitrile	9	9	0	0	9	9	9	0	0

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>			<u>C-S SYSTEMIC</u>			<u>G. COCKROACH</u>	
	<u>HOPPER</u>											
	ppm	100	1000	ppm	100	1000	ppm	7	days	100	100	1000
5-(3,4-dichloro-phenyl)-4-nitro-pyrrole-2-carbo-nitrile	9	9	9	9	9	9	9	9	9	0	0	9
2,4-dibromo-5-(p-chlorophenyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	9	9	9	9	0	0
3-bromo-5-[p-(tri-fluoromethoxy)phenyl]pyrrole-2,4-dicarbonitrile	9	9	9	9	9	9	9	9	9	9	0	9

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW³</u>		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>									
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES.
3-bromo-5-(3,4-dichlorophenyl)-4-nitropyrrole-2-carbonitrile	9	6	0	9	9	9	0	0	0	0
4-bromo-2-(p-chlorophenyl)-1-methyl-5-(trifluoromethyl)pyrrole-3-carbonitrile	9	9	0	9	9	9	0	0	0	9
3,4-dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile	0	9	9	9	9	9	0	0	0	9

Table IV (Cont..)

Compound	LEAF HOPPER		TBW ³		ARMYFORMS		C-S SYSTEMIC		G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYFORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	1000	ppm	100	ppm	1000	ppm	100	ppm	1000
2-(3,4-dichloro-phenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile	0	9	9	9	9	9	0	0	9	9
5-(trifluoromethyl)-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-3-carbonitrile	7	9	9	9	9	9	0	0	9	9

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBE³</u>		<u>ARTYFORMS</u>		<u>C-S SYSTEMIC</u>		<u>LEAF</u>		<u>G. COCKROACH</u>	
	<u>HOPPER</u>								<u>HOPPER</u>		<u>RES.</u>	
	ppm	100	1000	ppm	100	1000	ppm	100	ppm	100	ppm	1000
2,5-dibromo-4-(p-chlorophenyl)pyrrole-3-carbonitrile	0	9	0	9	9	9	0	0	0	0	9	
3,5-dibromo-4-(p-chlorophenyl)pyrrole-2-carbonitrile	0	9	3	9	9	9	0	0	0	0	0	
2-p-tolyl-5-(tri-fluoromethyl)pyrrole-3-carbonitrile	0	9	0	7.5	0	0	0	0	0	0	0	
4-bromo-2-p-tolyl-5-(trifluoromethyl)pyrrole-3-carbonitrile	0	9	6	9	9	9	-	-	-	-	0	

G. COCKROACH

113

Table IV (Cont.)

Compound	<u>LEAF</u>		<u>TBW</u> ³	<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>		
	<u>HOPPER</u>									
	ppm	1000	ppm	100	1000	ppm	7 days	SAW	LEAF HOPPER	RES.
	100	1000	ppm	100	1000	ppm	100	100	ppm	1000
5-(alpha, alpha, alpha-trifluoro-p-tolyl)pyrrole-2,4-dicarbonitrile	9	9	9	9	9	9	9	7	9	0
1-methyl-3-(alpha, alpha, alpha-trifluoro-p-tolyl)pyrrole-2,4-dicarbonitrile	0	3	0	0	9	0	0	0	0	0

Table IV (Cont.)

Compound	<u>LEAF</u> <u>HOPPER</u>		<u>TBW</u> ³		<u>ARMYWORMS</u>		<u>C-S SYSTEMIC</u>		<u>G. COCKROACH</u>	
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES.
4-bromo-5-(tri- fluoromethyl)-2- (alpha, alpha, alpha- trifluoro-p-tolyl) pyrrole-3-carbo- nitrile	9	9	9	9	9	9	0	0	0	9
3-bromo-1-methyl- 5-(alpha, alpha, alpha-trifluoro-p- tolyl)pyrrole-2,4- dicarbonitrile	0	7	0	9	0	9	-	-	-	0


Table IV (Cont.)

Compound	LEAF HOPPER		TBW ³		ARMYFORMS		C-S SYSTEMIC		G. COCKROACH	
	LEAF HOPPER		TBW ³		ARMYFORMS		C-S SYSTEMIC		G. COCKROACH	
	ppm	100	1000	ppm	100	1000	ppm	100	1000	RES. ppm
4,5-dichloro-2-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-1,3-dicarbonitrile	9	9	9	9	9	9	-	-	-	9
3-bromo-5-(alpha,alpha,alpha-trifluoro-p-tolyl)pyrrole-2,4-dicarbonitrile	-	-	-	-	-	-	-	-	-	-

Claims


Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. A compound having the formula I structure:



wherein X is F, Cl, Br, I, or CF₃; Y is F, Cl, Br, I, CF₃ or CN; W is CN or NO₂ and A is H; C₁-C₄ alkyl optionally substituted with from one to three halogen atoms, one hydroxy, one C₁-C₄ alkoxy or one C₁-C₄ alkylthio, one phenyl optionally substituted with C₁-C₃ alkyl or C₁-C₃ alkoxy or with one to three halogen atoms, one phenoxy optionally substituted with one to three halogen atoms or one benzyloxy optionally substituted with one halogen substituent; C₁-C₄ carbalkoxymethyl; C₃-C₄ alkenyl optionally substituted with from one to three halogen atoms; cyano; C₃-C₄ alkynyl optionally substituted with one halogen atom; di-(C₁-C₄ alkyl) aminocarbonyl; or C₄-C₆ cycloalkyl-aminocarbonyl; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:


-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCF₃, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2.


2. A compound according to claim 1 wherein A is hydrogen or C₁-C₄ alkoxyethyl; W is CN or NO₂; X and Y are each Cl, CF₃, or Br; R is F, Cl, Br, CF₃, or OCF₃; M is H, F, Cl, or Br; and L is H, or F.
3. A compound according to claim 1, 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile; 4,5-dichloro-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile; 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl)pyrrole; 2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole; 4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; 3,4-dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile; 2,4-dibromo-5-(p-chlorophenyl)pyrrole-3-carbonitrile; 5-(p-chlorophenyl)-3-(trifluoromethyl)pyrrole-2,4-dicarbonitrile; 3-bromo-5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonitrile; 4,5-dichloro-1-(ethoxymethyl)-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile; 4-bromo-2-(p-chloro-phenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; and 4-bromo-2-(3,4-dichlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile.
4. A method for controlling insects, nematodes and acarina comprising: contacting said insects, nematodes and acarina, their breeding grounds, food supply or habitat with an insecticidally, nematocidally

and acaricidally effective amount of a compound having the structure:



wherein X is H, F, Cl, Br, I, or CF₃; Y is F, Cl, Br, I, CF₃ or CN; W is CN or NO₂ and A is H; C₁-C₄ alkyl optionally substituted with from one to three halogen atoms, one hydroxy, one C₁-C₄ alkoxy or one C₁-C₄ alkylthio, one phenyl optionally substituted with C₁-C₃ alkyl or C₁-C₃ alkoxy or with one to three halogen atoms, one phenoxy optionally substituted with one to three halogen atoms or one benzyloxy optionally substituted with one halogen substituent; C₁-C₄ carbalkoxymethyl; C₃-C₄ alkenyl optionally substituted with from one to three halogen atoms; cyano; C₃-C₄ alkynyl optionally substituted with one halogen atom; di-(C₁-C₄ alkyl) aminocarbonyl; or C₄-C₆ cycloalkyl-aminocarbonyl; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:

-OCH₂O-, -OCF₂O- or




Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2.


5. A method according to claim 4 wherein said compound is:

- 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile;
 4,5-dichloro-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl) pyrrole;
 2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole;
 4-bromo-2-(p-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
 3,4-dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile;
 2,4-dibromo-5-(p-chlorophenyl)pyrrole-3-carbonitrile;
 5-(p-chlorophenyl)pyrrole-2,4-dicarbonitrile;
 3-bromo-5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonitrile;
 4,5-dichloro-1-(ethoxymethyl)-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile;
 4-bromo-2-(p-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; or
 4-bromo-2-(3,4-dichlorophenyl)-5-(trifluoromethyl) pyrrole-3-carbonitrile.

6. A method for protecting growing plants from attack by insects, nematodes and acarina, comprising applying to the foliage of said plants or to the soil or water in which they are growing, an insecticidally, nematocidally or acaricidally, effective amount of a formula I compound having the structure:




wherein X is H, F, Cl, Br, I, or CF₃; Y is F, Cl, Br, I, CF₃ or CN; W is CN or NO₂ and A is H; C₁-C₄ alkyl optionally substituted with from one to three halogen atoms, one hydroxy, one C₁-C₄ alkoxy or one C₁-C₄ alkylthio, one phenyl optionally substituted with C₁-C₃ alkyl or C₁-C₃ alkoxy or with one to three halogen atoms, one phenoxy optionally substituted with one to three halogen atoms or one benzyloxy optionally substituted with one halogen substituent; C₁-C₄ carbalkoxymethyl; C₃-C₄ alkenyl optionally substituted with from one to three halogen atoms; cyano; C₃-C₄ alkynyl optionally substituted with one halogen; di-(C₁-C₄ alkyl) aminocarbonyl; or C₄-C₆ cycloalkylaminocarbonyl; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:
-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2.


7. A method according to claim 6 wherein said compound is:
 - 4,5-dichloro-2-(3,4-dichlorophenyl)pyrrole-3-carbonitrile;
 - 4,5-dichloro-2-(α,α,α -trifluoro-*p*-tolyl)pyrrole-3-carbonitrile;
 - 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-*p*-tolyl) pyrrole;
 - 2,3-dichloro-5-(3,4-dichlorophenyl)-4-nitropyrrole;
 - 4-bromo-2-(*p*-chlorophenyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile;
 - 3,4-dibromo-5-(3,4-dichlorophenyl)pyrrole-2-carbonitrile;
 - 2,4-dibromo-5-(*p*-chlorophenyl)pyrrole-3-carbonitrile;
 - 5-(*p*-chlorophenyl)pyrrole-2,4-dicarbonitrile;
 - 3-bromo-5-(3,4-dichlorophenyl)pyrrole-2,4-dicarbonitrile;
 - 4,5-dichloro-1-(ethoxymethyl)-2-(α,α,α -trifluoro-*p*-tolyl)pyrrole-3-carbonitrile;
 - 4-bromo-2-(*p*-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)pyrrole-3-carbonitrile; or
 - 4-bromo-2-(3,4-dichlorophenyl)-5-(trifluoromethyl) pyrrole-3-carbonitrile.
8. A method according to claim 6 wherein said compound is applied to said plants or the soil in which they are growing, at about 0.125 kg/ha to about 4.0 kg/ha of said formula I compound.
9. A method according to claim 6 wherein said formula I compound is applied to the foliage of said plants or the soil or water in which they are growing, in the form of a liquid composition containing from about 10 ppm to about 10,000 ppm of said formula I compound.

10. A method for the preparation of a novel arylpyrrole compound having the structure:




wherein A, X and Y are as defined in claim 1; W is CN or NO₂; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:


-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃alkyl; and n is an integer of 0, 1 or 2; comprising, reacting a benzoylacetone nitrile or α-nitroacetophenone having the structure:




wherein L, M, R, and W are as described above with 2,2-di(C₁-C₄ alkoxy)ethylamine, at an elevated temperature, to yield an α-[2,2-di(C₁-C₄ alkoxy)ethylamino]-β-cyanostyrene or α-[2,2-di(C₁-C₄ alkoxy)-ethylamino]-β-nitrostyrene having the structure:



15 wherein L, M, R, and W are as described above and treating the thus formed α -[2,2-di(C₁-C₄ alkoxy)-ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene with a mineral or organic acid to yield the desired arylpyrrole.

11. A method according to claim 10 wherein the reaction of 2,2-di(C₁-C₄ alkoxy)ethylamine with benzoylacetoneitrile or α -nitroacetophenone is conducted neat or in the presence of an inert organic solvent.
- 20 12. A method according to claim 10 wherein the α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene is treated with hydrochloric, hydrobromic, or trifluoroacetic acid to form the arylpyrrole.
- 25 13. A process for the preparation of an α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene represented by the structure:




40 wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:

-OCH₂O-, -OCF₂O- or




50 Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCI, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2; comprising, reacting a benzoylacetoneitrile or α -nitroacetophenone having the structure:

55




wherein L, M, R, and W are as described above with 2,2-diethoxyethylamine, at an elevated temperature, to yield an α -(2,2-di(C₁-C₄ alkoxy)ethylamino)- β -cyanostyrene or α -(2,2-di(C₁-C₄ alkoxy)ethylamino)- β -nitrostyrene having the structure:




wherein L, M, R, and W are as described above.

14. A compound having the structural formula:



wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:

-OCH₂O-, -OCF₂O- or




Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2.

15. A compound according to claim 14 (E) p-chloro- β -[(formylmethyl)amino]cinnamitrile diethyl acetal; (Z) p-chloro- β -[(formylmethyl)amino]cinnamitrile diethyl acetal; β -[(formylmethyl)amino]-3,4-dimethoxycinnamitrile diethyl acetal; (Z)-methyl p-{2-cyano-1-[(formylmethyl)amino]vinyl}benzoic acid diethyl acetal; 3,4-dichloro- β [(formylmethyl)amino]cinnamitrile diethyl acetal; (Z)- β -[(formylmethyl)amino]-p-


methylcinnamonitrile diethyl acetal; β -[(formylmethyl)amino] *p*-trifluoromethoxycinnamonitrile dimethyl acetal; (E) *p*-chloro- β -[(formylmethyl)amino]cinnamonitrile dimethyl acetal; N-(formylmethyl)-*p*-methyl- α -(nitromethylene)benzylamine diethyl acetal; N-(formylmethyl)-3,4-dimethoxy- α -(nitromethylene)-benzylamine diethyl acetal; *p*-chloro-N-(formylmethyl)- α -(nitromethylene)benzylamine diethyl acetal; N-(formylmethyl)- α -(nitromethylene)-2-naphthenemethylamine diethyl acetal; methyl *p*-{ α -[(formylmethyl)amino]- β -nitrovinyl}benzoate *p*-(diethylacetal); and *p*-trifluoromethyl-N-[formylmethyl- α -(nitromethylene)-]benzylamine diethyl acetal.

16. A process for the preparation of a compound represented by the structure:




wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:


-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2; comprising, reacting a benzoylacetone nitrile or α -nitroacetophenone having the structure:




wherein L, M, R, and W are as described above with 2,2-di(C₁-C₄ alkoxy)ethylamine, at an elevated temperature, to give a compound having the structure:



wherein L, M, R, and W are as described above.


Claims for the following Contracting State : ES

1. A method for the preparation of a novel arylpyrrole compound having the structure




wherein X is F, Cl, Br, I, or CF₃; Y is F, Cl, Br, I, CF₃ or CN; W is CN or NO₂ and A is H; C₁-C₄ alkyl optionally substituted with from one to three halogen atoms, one hydroxy, one C₁-C₄ alkoxy or one C₁-C₄ alkylthio, one phenyl optionally substituted with C₁-C₃ alkyl or C₁-C₃ alkoxy or with one to three halogen atoms, one phenoxy optionally substituted with one to three halogen atoms or one benzyloxy optionally substituted with one halogen substituent; C₁-C₄ carbalkoxymethyl; C₃-C₄ alkenyl optionally substituted with from one to three halogen atoms; cyano; C₃-C₄ alkynyl optionally substituted with one halogen atom; di-(C₁-C₄ alkyl) aminocarbonyl; or C₄-C₆ cycloalkyl-aminocarbonyl; L is H, F, Cl or Br; and M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO or NR₃R₄, and when M and R are on adjacent positions and taken with the carbon atoms to which they are attached they may form a ring in which MR represents the structure:


-OCH₂O-, -OCF₂O- or



Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2 comprising reacting a benzoyl acetonitrile or α-nitroacetophenone having the structure:




wherein L, M, R, and W are as described above with 2,2-di(C₁-C₄ alkoxy)ethylamine, at an elevated temperature, to yield an α-[2,2-di(C₁-C₄ alkoxy)ethylamino]-β-cyanostyrene or α-[2,2-di(C₁-C₄ alkoxy)-ethylamino]-β-nitrostyrene having the structure:



wherein L, M, R, and W are as described above and treating the thus formed α -[2,2-di(C₁-C₄ alkoxy)-ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)amino]- β -nitrostyrene with a mineral or organic acid to yield the desired arylpyrrole.

15

2. A method according to claim 1 wherein the reaction of 2,2-di(C₁-C₄ alkoxy)ethylamine with benzoylacetonitrile or α -nitroacetophenone is conducted neat or in the presence of an inert organic solvent.
 - 20 3. A method according to claim 1 wherein the α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene is treated with hydrochloric, hydrobromic, or trifluoroacetic acid to form the arylpyrrole.
 4. A process for the preparation of an α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -cyanostyrene or α -[2,2-di(C₁-C₄ alkoxy)ethylamino]- β -nitrostyrene represented by the structure:
- 25



wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:


40

-OCH₂O-, -OCF₂O- or




50 Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2; comprising, reacting a benzoylacetonitrile or α -nitroacetophenone having the structure:

55




10 wherein L, M, R, and W are as described above with 2,2-diethoxyethylamine, at an elevated temperature, to yield an α -(2,2-di(C₁-C₄ alkoxy)ethylamino)- β -cyanostyrene or α -(2,2-di(C₁-C₄ alkoxy)ethylamino)- β -nitrostyrene having the structure:




25 wherein L, M, R, and W are as described above.

5. A process for the preparation of a compound represented by the structure:




40 wherein W is CN or NO₂; L is H, F, Cl, or Br; M and R are each independently H, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, C₁-C₃ alkylsulfinyl, C₁-C₃ alkylsulfonyl, cyano, F, Cl, Br, I, nitro, CF₃, R₁CF₂Z, R₂CO, or NR₃R₄ and when on adjacent positions and taken together with the carbon atoms to which they are attached M and R may form a ring in which MR represent the structure:

-OCH₂O-, -OCF₂O- or




50 Z is S(O)_n or O; R₁ is H, F, CHF₂, CHFCl, or CF₃; R₂ is C₁-C₃ alkyl, C₁-C₃ alkoxy, or NR₃R₄; R₃ is H or C₁-C₃ alkyl; R₄ is H, C₁-C₃ alkyl, or R₅CO; R₅ is H or C₁-C₃ alkyl; and n is an integer of 0, 1 or 2; comprising, reacting a benzoylacetonitrile or α -nitroacetophenone having the structure:



5

10 wherein L, M, R, and W are as described above with 2,2-di(C₁-C₄ alkoxy)ethylamine, at an elevated temperature, to give a compound having the structure:



15


20

wherein L, M, R, and W are as described above.

Patentansprüche

25 Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. Verbindung der Formel (I):



30


35

40

45 worin X F, Cl, Br, I oder CF₃ ist; Y F, Cl, Br, I, CF₃ oder CN ist; W CN oder NO₂ ist und A H; C₁-C₄-Alkyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen, einem Hydroxy, einem C₁-C₄-Alkoxy oder einem C₁-C₄-Alkylthio, einem Phenyl, gegebenenfalls substituiert mit C₁-C₃-Alkyl oder C₁-C₃-Alkoxy oder mit ein bis drei Halogenatomen, einem Phenoxy, gegebenenfalls substituiert mit ein bis drei Halogenatomen oder einem Benzyloxy, gegebenenfalls substituiert mit einem Halogenatom; C₁-C₄-Carbalkoxymethyl; C₃-C₄-Alkenyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen; Cyan; C₃-C₄-Alkynyl, gegebenenfalls substituiert mit einem Halogenatom; Di(C₁-C₄-alkyl)aminocarbonyl; oder C₄-C₆-Cycloalkylaminocarbonyl ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

55

-OCH₂O-, -OCF₂O- oder




Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist.

2. Verbindung nach Anspruch 1, worin A Wasserstoff oder C₁-C₄-Alkoxymethyl ist; W CN oder NO₂ ist; X und Y jeweils Cl, CF₃ oder Br sind; R F, Cl, Br, CF₃ oder OCF₃ ist; M H, F, Cl oder Br ist und L H oder F ist.


3. Verbindung nach Anspruch 1, 4,5-Dichlor-2-(3,4-dichlorphenyl)pyrrol-3-carbonitril; 4,5-Dichlor-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril; 2,3-Dichlor-4-nitro-5-(α,α,α -trifluor-p-tolyl)pyrrol; 2,3-Dichlor-5-(3,4-dichlorphenyl)-4-nitropyrrol; 4-Brom-2-(p-chlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril; 3,4-Dibrom-5-(3,4-dichlorphenyl)pyrrol-2-carbonitril; 2,4-Dibrom-5-(p-chlorphenyl)pyrrol-3-carbonitril; 5-(p-Chlorphenyl)-3-(trifluormethyl)pyrrol-2,4-dicarbonitril; 3-Brom-5-(3,4-dichlorphenyl)pyrrol-2,4-dicarbonitril; 4,5-Dichlor-1-(ethoxymethyl)-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril; 4-Brom-2-(p-chlorphenyl)-1-(ethoxymethyl)-5-(trifluormethyl)pyrrol-3-carbonitril und 4-Brom-2-(3,4-dichlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril.

4. Verfahren zum Bekämpfen von Insekten, Nematoden und Acarina, umfassend:
In Berührung bringen der Insekten, Nematoden und Acarina, ihre Brutstätten, Nahrungszufuhr oder Wohnstätte mit einer insektizid, nematizid und acarizid wirksamen Menge einer Verbindung mit der Struktur



worin X F, Cl, Br, I oder CF₃ ist; Y F, Cl, Br, I, CF₃ oder CN ist; W CN oder NO₂ ist und A H; C₁-C₄-Alkyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen, einem Hydroxy, einem C₁-C₄-Alkoxy oder einem C₁-C₄-Alkylthio, einem Phenyl, gegebenenfalls substituiert mit C₁-C₃-Alkyl oder C₁-C₃-Alkoxy oder mit ein bis drei Halogenatomen, einem Phenoxy, gegebenenfalls substituiert mit ein bis drei Halogenatomen oder einem Benzyloxy, gegebenenfalls substituiert mit einem Halogenatom; C₁-C₄-Carbalkoxymethyl; C₃-C₄-Alkenyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen; Cyan; C₃-C₄-Alkynyl, gegebenenfalls substituiert mit einem Halogenatom; Di(C₁-C₄-alkyl)aminocarbonyl; oder C₄-C₆-Cycloalkylaminocarbonyl ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:


-OCH₂O-, -OCF₃O- oder



5

10 Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist.

5. Verfahren nach Anspruch 4, worin die Verbindung ist:
- 15 4,5-Dichlor-2-(3,4-di-chlorphenyl)pyrrol-3-carbonitril;
4,5-Dichlor-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril;
2,3-Dichlor-4-nitro-5-(α,α,α -trifluor-p-tolyl)pyrrol;
2,3-Dichlor-5-(3,4-dichlorphenyl)-4-nitropyrrol;
4-Brom-2-(p-chlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril;
3,4-Dibrom-5-(3,4-dichlorphenyl)pyrrol-2-carbonitril;
20 2,4-Dibrom-5-(p-chlorphenyl)pyrrol-3-carbonitril;
5-(p-Chlorphenyl)-3-(trifluormethyl)pyrrol-2,4-dicarbonitril;
3-Brom-5-(3,4-dichlorphenyl)pyrrol-2,4-dicarbonitril;
4,5-Dichlor-1-(ethoxymethyl)-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril;
4-Brom-2-(p-chlor-phenyl)-1-(ethoxymethyl)-5-(trifluormethyl)pyrrol-3-carbonitril oder
25 4-Brom-2-(3,4-dichlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril.
6. Verfahren zum Schützen wachsender Pflanzen gegen Angriff durch Insekten, Nematoden und Acarina, umfassend das Aufbringen auf das Laub der Pflanzen oder den Boden oder das Wasser, in dem sie wachsen, einer insektizid, nematizid oder acarizid wirksamen Menge einer Verbindung der Formel (I) mit der Struktur:
- 30



35


40

worin X F, Cl, Br, I oder CF₃ ist; Y F, Cl, Br, I, CF₃ oder CN ist; W CN oder NO₂ ist und A H; C₁-C₄-Alkyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen, einem Hydroxy, einem C₁-C₄-Alkoxy oder einem C₁-C₄-Alkylthio, einem Phenyl, gegebenenfalls substituiert mit C₁-C₃-Alkyl oder C₁-C₃-Alkoxy oder mit ein bis drei Halogenatomen, einem Phenoxy, gegebenenfalls substituiert mit ein bis drei Halogenatomen oder einem Benzyloxy, gegebenenfalls substituiert mit einem Halogenatom; C₁-C₄-Carbalkoxymethyl; C₃-C₄-Alkenyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen; Cyan; C₃-C₄-Alkynyl, gegebenenfalls substituiert mit einem Halogenatom; Di(C₁-C₄-alkyl)aminocarbonyl; oder C₄-C₆-Cycloalkylaminocarbonyl ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

50

-OCH₂O-, -OCF₂O- oder

55




5

10 Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist.

7. Verfahren nach Anspruch 6, worin die Verbindung ist:
- 15 4,5-Dichlor-2-(3,4-di-chlorphenyl)pyrrol-3-carbonitril;
 4,5-Dichlor-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril;
 2,3-Dichlor-4-nitro-5-(α,α,α -trifluor-p-tolyl)pyrrol;
 2,3-Dichlor-5-(3,4-dichlorphenyl)-4-nitropyrrol;
 4-Brom-2-(p-chlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril;
 3,4-Dibrom-5-(3,4-dichlorphenyl)pyrrol-2-carbonitril;
 20 2,4-Dibrom-5-(p-chlorphenyl)pyrrol-3-carbonitril;
 5-(p-Chlorphenyl)-3-(trifluormethyl)pyrrol-2,4-dicarbonitril;
 3-Brom-5-(3,4-dichlorphenyl)pyrrol-2,4-dicarbonitril;
 4,5-Dichlor-1-(ethoxymethyl)-2-(α,α,α -trifluor-p-tolyl)pyrrol-3-carbonitril;
 4-Brom-2-(p-chlor-phenyl)-1-(ethoxymethyl)-5-(trifluormethyl)pyrrol-3-carbonitril oder
 25 4-Brom-2-(3,4-dichlorphenyl)-5-(trifluormethyl)pyrrol-3-carbonitril.
8. Verfahren nach Anspruch 6, worin die Verbindung auf die Pflanzen oder den Boden, in dem sie wachsen in einer Menge von etwa 0,125 kg/ha bis etwa 4,0 kg/ha der Verbindung der Formel I aufgebracht wird.
- 30 9. Verfahren nach Anspruch 6, worin die Verbindung der Formel I auf das Laub der Pflanzen oder den Boden oder das Wasser, in dem sie wachsen, in Form einer flüssigen Zusammensetzung aufgebracht wird, die von etwa 10 ppm bis etwa 10.000 ppm der Verbindung der Formel I enthält.
- 35 10. Verfahren zum Herstellen einer neuen Arylpyrrol-Verbindung mit der Struktur:

40




45

50 worin A, X und Y die in Anspruch 1 genannte Bedeutung haben; W ist CN oder NO₂; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

-OCH₂O-, -OCF₂O- oder

55




5

10 Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist, umfassend das Umsetzen eines Benzoylacetonnitrils oder α-Nitroacetophenons mit der Struktur:

15

20




25

worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Di(C₁-C₄-alkoxy)ethylamin bei einer erhöhten Temperatur, um ein α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-cyanstyrol oder α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-nitrostyrol mit der Struktur:

30

35



40

worin L, M, R und W die oben genannte Bedeutung haben, zu erhalten und Behandeln des so gebildeten α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-cyanstyrols oder α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-nitrostyrols mit einer Mineral- oder organischen Säure, um das erwünschte Arylpyrrol zu ergeben.

45


11. Verfahren nach Anspruch 10, worin die Umsetzung von 2,2-Di(C₁-C₄-alkoxy)ethylamin mit Benzoylacetonnitril oder α-Nitroacetophenon unverdünnt oder in Gegenwart eines inerten organischen Lösungsmittels ausgeführt wird.

50

12. Verfahren nach Anspruch 10, worin das α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-cyanstyrol oder α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-nitrostyrol mit Chlowsasserstoff-, Bromwasserstoff- oder Trifluoressigsäure behandelt wird, um das Arylpyrrol zu bilden.


13. Verfahren zum Herstellen eines α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-cyanstyrols oder α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-nitrostyrols, dargestellt durch die Struktur:

55




15 worin W CN oder NO₂ ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:


-OCH₂O-, -OCF₂O- oder



30 Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCl oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist, umfassend das Umsetzen eines Benzoylacetonnitrils oder α-Nitroacetophenons der Struktur:



45 worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Diethoxyethylamin bei einer erhöhten Temperatur, um ein α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-cyanstyrol oder α-[2,2-Di(C₁-C₄-alkoxy)ethylamino]-β-nitrostyrol mit der Struktur:




worin L, M, R und W die oben genannte Bedeutung haben, herzustellen.

14. Verbindung mit der Strukturformel:

5

10




15

worin W CN oder NO₂ ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

-OCH₂O-, -OCF₂O- oder

20

25



Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist.

30

15. Verbindung nach Anspruch 14,


(E) p-Chlor-β-[(formylmethyl)amino]-zimtsäurenitril-diethyl-acetal; (Z) p-Chlor-β-[(formylmethyl)amino]-zimtsäurenitril-diethyl-acetal; β-[(Formylmethyl)amino]-3,4-dimethoxyzimtsäurenitril-diethyl-acetal; (Z)-Methyl-p-(2-cyan-1-[(formylmethyl)amino]vinyl)benzoesäure-diethylacetal; 3,4-Dichlor-β-[(formylmethyl)amino]-zimtsäurenitril-diethyl-acetal; (Z)-β-[(formylmethyl)amino]-p-methylzimtsäurenitril-diethyl-acetal; β-[(Formylmethyl)amino]-p-trifluormethoxyzimtsäurenitril-dimethylacetal; (E) p-Chlor-β-[(formylmethyl)amino]-zimtsäurenitril-dimethylacetal; N-(Formylmethyl)-p-methyl-α-(nitromethylen)-benzylamin-diethylacetal, N-(Formylmethyl)-3,4-dimethoxy-α-(nitromethylen)benzylamin-diethylacetal, p-Chlor-N-(formylmethyl)-α-(nitromethylen)benzylamin-diethylacetal; N-(Formylmethyl)-α-(nitromethylen)-2-naphthenmethyamin-diethylacetal; Methyl-p-α[(formylmethyl)amino]-β-nitrovinylbenzoat-p-(diethylacetal) und p-Trifluormethyl-N-[formylmethyl-α-(nitromethylen)]benzylamin-diethylacetal.

40

16. Verfahren zum Herstellen einer Verbindung, die durch die Struktur repräsentiert ist:

45

50




55

worin W CN oder NO₂ ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit


den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

-OCH₂O-, -OCF₂O- oder




Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCl oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist, umfassend

Umsetzen eines Benzoylacetonitrils oder α-Nitroacetophenons der Struktur:




worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Di(C₁-C₄-alkoxy)ethylamin bei einer erhöhten Temperatur, um eine Verbindung mit der Struktur:



zu erhalten, worin L, M, R und W die oben genannte Bedeutung haben.


Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zum Herstellen einer neuen Arylpyrrol-Verbindung mit der Struktur




worin X F, Cl, Br, I oder CF₃ ist; Y F, Cl, Br, I, CF₃ oder CN ist; W CN oder NO₂ ist und A H; C₁-C₄-Alkyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen, einem Hydroxy, einem C₁-C₄-Alkoxy oder einem C₁-C₄-Alkylthio, einem Phenyl, gegebenenfalls substituiert mit C₁-C₃-Alkyl oder C₁-C₃-Alkoxy oder mit ein bis drei Halogenatomen, einem Phenoxy, gegebenenfalls substituiert mit ein bis drei Halogenatomen oder einem Benzyloxy, gegebenenfalls substituiert mit einem Halogenatom; C₁-C₄-Carbalkoxymethyl; C₃-C₄-Alkenyl, gegebenenfalls substituiert mit ein bis drei Halogenatomen; Cyan; C₃-C₄-Alkynyl, gegebenenfalls substituiert mit einem Halogenatom; Di(C₁-C₄-alkyl)aminocarbonyl; oder C₄-C₆-Cycloalkylaminocarbonyl ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfinyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:


-OCH₂O-, OCF₂O- oder



Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine ganze Zahl aus 0, 1 oder 2 ist, umfassend das Umsetzen eines Benzoylacetonitrils oder α-Nitroacetophenons mit der Struktur:



worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Di(C₁-C₄-alkoxy)ethylamin bei einer erhöhten Temperatur, um ein α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -cyanstyrol oder α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrol mit der Struktur:




15 worin L, M, R und W die oben genannte Bedeutung haben, zu erhalten und Behandeln des so gebildeten α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -cyanstyrols oder α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrols mit einer Mineral- oder organischen Säure, um das erwünschte Arylpyrrol zu ergeben.

20 2. Verfahren nach Anspruch 1, worin die Umsetzung von 2,2-Di(C₁-C₄-alkoxy)ethylamin mit Benzoylacetonitril oder α -Nitroacetophenon unverdünnt oder in Gegenwart eines inerten organischen Lösungsmittels ausgeführt wird.

25 3. Verfahren nach Anspruch 1, worin das α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -cyanstyrol oder α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrol mit Chlowsasserstoff-, Bromwasserstoff- oder Trifluoressigsäure behandelt wird, um das Arylpyrrol zu bilden.

30 4. Verfahren zum Herstellen eines α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -cyanstyrols oder α -(2,2-Di(C₁-C₄-alkoxy)ethylamino)- β -nitrostyrols, dargestellt durch die Struktur:




45 worin W CN oder NO₂ ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:

-OCH₂O-, -OCF₂O- oder




Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCl oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine

ganze Zahl aus 0, 1 oder 2 ist, umfassend das Umsetzen eines Benzoylacetonnitrils oder α -Nitroacetophenons der Struktur:




worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Diethoxyethylamin bei einer erhöhten Temperatur, um ein α -[2,2-Di(C₁-C₄-alkoxy)ethylamino]- β -cyanstyrol oder α -[2,2-Di(C₁-C₄-alkoxy)ethylamino]- β -nitrostyrol mit der Struktur:




worin L, M, R und W die oben genannte Bedeutung haben, zu ergeben.

5. Verfahren zum Herstellen einer Verbindung, die durch die Struktur repräsentiert ist:



worin W CN oder NO₂ ist; L H, F, Cl oder Br ist; und M und R jeweils unabhängig H, C₁-C₃-Alkyl, C₁-C₃-Alkoxy, C₁-C₃-Alkylthio, C₁-C₃-Alkylsulfonyl, C₁-C₃-Alkylsulfonyl, Cyan, F, Cl, Br, I, Nitro, CF₃, R₁CF₂Z, R₂CO oder N₃R₄ sind und, wenn M und R sich in benachbarten Positionen befinden und mit den Kohlenstoffatomen, an die sie gebunden sind, zusammen genommen werden, sie einen Ring bilden können, in dem MR die Struktur repräsentiert:


-OCH₂O-, -OCF₂O- oder




Z S(O)_n oder O ist; R₁ H, F, CHF₂, CHFCI oder CF₃ ist; R₂ C₁-C₃-Alkyl, C₁-C₃-Alkoxy oder NR₃R₄ ist; R₃ H oder C₁-C₃-Alkyl ist; R₄ H, C₁-C₃-Alkyl oder R₅CO ist; R₅ H oder C₁-C₃-Alkyl ist und n eine

ganze Zahl aus 0, 1 oder 2 ist, umfassend

Umsetzen eines Benzoylacetonnitrils oder α -Nitroacetophenons der Struktur:



worin L, M, R und W die oben genannte Bedeutung haben, mit 2,2-Di(C₁-C₄-alkoxy)ethylamin bei einer erhöhten Temperatur, um eine Verbindung mit der Struktur:




zu erhalten, worin L, M, R und W die oben genannte Bedeutung haben.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE


1. Composé répondant à la formule développée I :



dans laquelle X est F, Cl, Br, I ou CF₃ ; Y est F, Cl, Br, I, CF₃ ou CN ; W est CN ou NO₂ et A est H ; un groupe alkyle en C₁-C₄ facultativement substitué avec un à trois atomes d'halogène, un groupe hydroxy, un groupe alcoxy en C₁-C₄ ou un groupe alkylthio en C₁-C₄, un groupe phényle facultativement substitué avec un groupe alkyle en C₁-C₃ ou un groupe alcoxy en C₁-C₃ ou avec un à trois atomes d'halogène, un groupe phénoxy facultativement substitué avec un à trois atomes d'halogène ou un groupe benzyloxy facultativement substitué avec un substituant halogène ; un groupe (carbalcoxy en C₁-C₄)méthyle ; un groupe alcényle en C₃-C₄ facultativement substitué avec un à trois atomes d'halogène ; un groupe cyano ; un groupe alcynyle en C₃-C₄ facultativement substitué avec un atome d'halogène ; un groupe di(alkyl en C₁-C₄) aminocarbonyle ; ou un groupe cycloalkyl(en C₁-C₄)-aminocarbonyle ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en

C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsque M et R sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, ils peuvent former un cycle dans lequel MR représente la structure :

-OCH₂O-, -OCF₂O- ou




Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2.

2. Composé selon la revendication 1, où A est un atome d'hydrogène ou un groupe (alcoxy en C₁-C₄)-méthyle ; W est CN ou NO₂ ; X et Y sont chacun Cl, CF₃ ou Br ; R est F, Cl, Br, CF₃ ou OCF₃ ; M est H, F, Cl ou Br ; et L est H ou F.


3. Composé selon la revendication 1 qui est le 4,5-dichloro-2-(3,4-dichlorophényl)pyrrole-3-carbonitrile ; le 4,5-dichloro-2-(α,α,α-trifluoro-p-tolyl)pyrrole-3-carbonitrile ; le 2,3-dichloro-4-nitro-5-(α,α,α-trifluoro-p-tolyl)pyrrole ; le 2,3-dichloro-5-(3,4-dichlorophényl)-4-nitropyrrole ; le 4-bromo-2-(p-chlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile ; le 3,4-dibromo-5-(3,4-dichlorophényl)pyrrole-2-carbonitrile ; le 2,4-dibromo-5-(p-chlorophényl)pyrrole-3-carbonitrile ; le 5-(p-chlorophényl)-3-(trifluorométhyl)pyrrole-2,4-dicarbonitrile ; le 3-bromo-5-(3,4-dichlorophényl)pyrrole-2,4-dicarbonitrile ; le 4,5-dichloro-1-(éthoxyméthyl)-2-(α,α,α-trifluoro-p-tolyl)pyrrole-3-carbonitrile ; le 4-bromo-2-(p-chlorophényl)-1-(éthoxyméthyl)-5-(trifluorométhyl)pyrrole-3-carbonitrile ; ou le 4-bromo-2-(3,4-dichlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile.

4. Procédé de maîtrise des insectes, des nématodes et des acariens comprenant : le contact desdits insectes, nématodes et acariens, de leur base de reproduction, de leur alimentation ou de leur habitat avec une quantité insecticide, nématicide et acaricide d'un composé de structure :



dans laquelle X est H, F, Cl, Br, I ou CF₃ ; Y est F, Cl, Br, I, CF₃ ou CN ; W est CN ou NO₂ et A est H ; un groupe alkyle en C₁-C₄ facultativement substitué avec un à trois atomes d'halogène, un groupe hydroxy, un groupe alcoxy en C₁-C₄ ou un groupe alkylthio en C₁-C₄, un groupe phényle facultativement substitué avec un groupe alkyle en C₁-C₃ ou un groupe alcoxy en C₁-C₃ ou avec un à trois atomes d'halogène, un groupe phénoxy facultativement substitué avec un à trois atomes d'halogène ou un groupe benzyloxy facultativement substitué avec un substituant halogène ; un groupe (carbalcoxy en C₁-C₄)-méthyle ; un groupe alcényle en C₃-C₄ facultativement substitué avec un à trois atomes d'halogène ; un groupe cyano ; un groupe alcynyle en C₃-C₄ facultativement substitué avec un atome d'halogène ; un groupe di(alkyl en C₁-C₄) aminocarbonyl ; ou un groupe cycloalkyl(en C₄-C₆)-aminocarbonyl ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsque M et R sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, ils peuvent former un cycle dans lequel MR représente la structure :

-OCH₂O-, -OCF₂O- ou




Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2.

5. Procédé selon la revendication 4, dans lequel ledit composé est :


- le 4,5-dichloro-2-(3,4-dichlorophényl)pyrrole-3-carbonitrile ;
- le 4,5-dichloro-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile ;
- le 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl)pyrrole ;
- le 2,3-dichloro-5-(3,4-dichlorophényl)-4-nitropyrrole ;
- le 4-bromo-2-(p-chlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile ;
- le 3,4-dibromo-5-(3,4-dichlorophényl)pyrrole-2-carbonitrile ;
- le 2,4-dibromo-5-(p-chlorophényl)pyrrole-3-carbonitrile ;
- le 5-(p-chlorophényl)pyrrole-2,4-dicarbonitrile ;
- le 3-bromo-5-(3,4-dichlorophényl)pyrrole-2,4-dicarbonitrile ;
- le 4,5-dichloro-1-(éthoxyméthyl)-2-(α,α,α -trifluoro-p-tolyl) pyrrole-3-carbonitrile ;
- le 4-bromo-2-(p-chlorophényl)-1-(éthoxyméthyl)-5-(trifluorométhyl) pyrrole-3-carbonitrile ; ou
- le 4-bromo-2-(3,4-dichlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile.

6. Procédé pour protéger les plantes en cours de croissance contre l'attaque par les insectes, les nématodes et les acariens, comprenant l'application, au feuillage desdites plantes ou au sol ou à l'eau où elles se développent, d'une quantité insecticide, nématocide ou acaricide efficace d'un composé de formule I ayant la structure :



dans laquelle X est H, F, Cl, Br, I ou CF₃ ; Y est F, Cl, Br, I, CF₃ ou CN ; W est CN ou NO₂ et A est H ; un groupe alkyle en C₁-C₄ facultativement substitué avec un à trois atomes d'halogène, un groupe hydroxy, un groupe alcoxy en C₁-C₄ ou un groupe alkylthio en C₁-C₄, un groupe phényle facultativement substitué avec un groupe alkyle en C₁-C₃ ou un groupe alcoxy en C₁-C₃ ou avec un à trois atomes d'halogène, un groupe phénoxy facultativement substitué avec un à trois atomes d'halogène ou un groupe benzyloxy facultativement substitué avec un substituant halogène ; un groupe (carbalcoxy en C₁-C₄)méthyle ; un groupe alcényle en C₃-C₄ facultativement substitué avec un à trois atomes d'halogène ; un groupe cyano ; un groupe alcynyle en C₃-C₄ facultativement substitué avec un atome d'halogène ; un groupe di(alkyl en C₁-C₄) aminocarbonyl ; ou un groupe cycloalkyl(en C₁-C₄)aminocarbonyl ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsque M et R sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, ils peuvent former un cycle dans lequel MR représente la structure :

-OCH₂O-, -OCF₂O- ou




5

10 Z est $S(O)_n$ ou O ; R_1 est H, F, CHF_2 , CH_2Cl ou CF_3 ; R_2 est un groupe alkyle en C_1-C_3 , un groupe alcoxy en C_1-C_3 ou NR_3R_4 ; R_3 est H ou un groupe alkyle en C_1-C_3 ; R_4 est H, un groupe alkyle en C_1-C_3 ou R_5CO ; R_5 est H ou un groupe alkyle en C_1-C_3 ; et n est un entier valant 0, 1 ou 2.

7. Procédé selon la revendication 6, dans lequel ledit composé est :
- le 4,5-dichloro-2-(3,4-dichlorophényl)pyrrole-3-carbonitrile ;
 - 15 le 4,5-dichloro-2-(α,α,α -trifluoro-p-tolyl)pyrrole-3-carbonitrile ;
 - le 2,3-dichloro-4-nitro-5-(α,α,α -trifluoro-p-tolyl)pyrrole ;
 - le 2,3-dichloro-5-(3,4-dichlorophényl)-4-nitropyrrole ;
 - le 4-bromo-2-(p-chlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile ;
 - le 3,4-dibromo-5-(3,4-dichlorophényl)pyrrole-2-carbonitrile ;
 - 20 le 2,4-dibromo-5-(p-chlorophényl)pyrrole-3-carbonitrile ;
 - le 5-(p-chlorophényl)pyrrole-2,4-dicarbonitrile ;
 - le 3-bromo-5-(3,4-dichlorophényl)pyrrole-2,4-dicarbonitrile ;
 - le 4,5-dichloro-1-(éthoxyméthyl)-2-(α,α,α -trifluoro-p-tolyl) pyrrole-3-carbonitrile ;
 - le 4-bromo-2-(p-chlorophényl)-1-(éthoxyméthyl)-5-(trifluorométhyl) pyrrole-3-carbonitrile ; ou
 - 25 le 4-bromo-2-(3,4-dichlorophényl)-5-(trifluorométhyl)pyrrole-3-carbonitrile.
8. Procédé selon la revendication 6, dans lequel ledit composé est appliqué auxdites plantes ou au sol où elles se développent, à raison d'environ 0,125 kg/ha à environ 4,0 kg/ha dudit composé de formule I.
- 30 9. Procédé selon la revendication 6, dans lequel on applique ledit composé de formule I au feuillage desdites plantes ou au sol ou à l'eau où elles se développent, sous forme d'une composition liquide contenant environ 10 ppm à environ 10 000 ppm dudit composé de formule I.

10. Procédé pour la préparation d'un nouvel arylpyrrole ayant la structure :

35




40


45

dans laquelle A, X et Y sont tels que définis à la revendication 1, W est CN ou NO_2 ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C_1-C_3 , un groupe alcoxy en C_1-C_3 , un groupe alkylthio en C_1-C_3 , un groupe alkylsulfinyle en C_1-C_3 , un groupe alkylsulfonyle en C_1-C_3 , un groupe cyano, F, Cl, Br, I, un groupe nitro, CF_3 , R_1CF_2Z , R_2CO ou NR_3R_4 et, lorsque M et R sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, ils peuvent former un cycle dans lequel MR représente la structure :


55



-OCH₂O-, -OCF₂O- ou Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α-nitroacétophénone ayant la structure :




dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-di(alcoxy en C₁-C₄)éthylamine, à une température élevée, pour former un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène ayant la structure :




dans laquelle L, M, R et W sont décrits comme ci-dessus, et le traitement de l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou de l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène ainsi formé avec un acide minéral ou organique pour former l'arylpyrrole désiré.


11. Procédé selon la revendication 10, dans lequel la réaction de la 2,2-di(alcoxy en C₁-C₄)éthylamine avec le benzoylacétonitrile ou l'α-nitroacétophénone est effectuée sans solvant ou en présence d'un solvant organique inerte.
12. Procédé selon la revendication 10, dans lequel on traite l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène avec l'acide chlorhydrique, bromhydrique ou trifluoroacétique pour former l'arylpyrrole.
13. Procédé pour la préparation d'un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou d'un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène, représenté par la structure :




15 dans laquelle W est CN ou NO₂ ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsqu'ils sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, M et R peuvent former un cycle dans lequel MR représente la structure :



25 Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α-nitroacétophénone ayant la structure :




45 dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-diéthoxyéthylamine, à une température élevée, pour former un α-(2,2-di(alcoxy en C₁-C₄)éthylamino)-β-cyanostyrène ou un α-(2,2-di(alcoxy en C₁-C₄)éthylamino)-β-nitrostyrène ayant la structure :



dans laquelle L, M, R et W sont décrits comme ci-dessus.

14. Composé répondant à la formule développée :



dans laquelle W est CN ou NO₂, L est H, F, Cl ou Br ; M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsqu'ils sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, M et R peuvent former un cycle dans lequel MR représente la structure :


-OCH₂O-, -OCF₂O- ou



25 Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2.


15. Composé selon la revendication 14 qui est le (E)-p-chloro-β-[(formylméthyl)amino]cinnamitrile-diéthylacétal ; le (Z)-p-chloro-β-[(formylméthyl)amino]cinnamitrile-diéthylacétal ; le β-[(formylméthyl)amino]-3,4-diméthoxycinnamitrile-diéthylacétal ; le (Z)-p-[2-cyano-1-[(formylméthyl)amino]vinyl]benzoate de méthyle-diéthylacétal ; le 3,4-dichloro-β-[(formylméthyl)amino]cinnamitrile-diéthylacétal ; le (Z)-β-[(formylméthyl)amino]-p-méthylcinnamitrile-diéthylacétal ; le β-[(formylméthyl)amino]-p-trifluorométhoxycinnamitrile-diéthylacétal ; le (E)-p-chloro-β-[(formylméthyl)amino]cinnamitrile-diéthylacétal ; le N-(formylméthyl)-p-méthyl-α-(nitrométhylène)benzylamine-diéthylacétal ; le N-(formylméthyl)-3,4-diméthoxy-α-(nitrométhylène)benzylamine-diéthylacétal ; le p-chloro-N-(formylméthyl)-α-(nitrométhylène)benzylamine-diéthylacétal ; le N-(formylméthyl)-α-(nitrométhylène)-2-naphtène-méthylamine-diéthylacétal ; le p-α-[(formylméthyl)amino]-β-nitrovinyl]benzoate de méthyle-p-diéthylacétal ; ou le p-trifluorométhyl-N-[(formylméthyl)-α-(nitrométhylène)]benzylamine-diéthylacétal.

16. Procédé pour la préparation d'un composé représenté par la structure :



dans laquelle W est CN ou NO₂, L est H, F, Cl ou Br ; M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsqu'ils sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, M et R peuvent former un cycle dans lequel MR représente la structure :


-OCH₂O-, -OCF₂O- ou



5

Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α-nitroacétophénone ayant la structure :


15



20

dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-di(alcoxy en C₁-C₄)éthylamine, à une température élevée, pour obtenir un composé ayant la structure :

25



30


dans laquelle L, M, R et W sont décrits comme ci-dessus.

35

Revendications pour l'Etat contractant suivant : ES

1. Procédé pour la préparation d'un nouveau dérivé d'arylpyrrole de formule

40



45


50

dans laquelle X est F, Cl, Br, I ou CF₃ ; Y est F, Cl, Br, I, CF₃ ou CN ; W est CN ou NO₂ et A est H : un groupe alkyle en C₁-C₄ facultativement substitué avec un à trois atomes d'halogène, un groupe hydroxy, un groupe alcoxy en C₁-C₄ ou un groupe alkylthio en C₁-C₄, un groupe phényle facultativement substitué avec un groupe alkyle en C₁-C₃ ou un groupe alcoxy en C₁-C₃ ou avec un à trois


55

atomes d'halogène, un groupe phénoxy facultativement substitué avec un à trois atomes d'halogène ou un groupe benzyloxy facultativement substitué avec un substituant halogène ; un groupe (carbalcoxy en C₁-C₄) méthyle ; un groupe alcényle en C₃-C₄ facultativement substitué avec un à trois atomes d'halogène ; un groupe cyano ; un groupe alcynyle en C₃-C₄ facultativement substitué avec un atome d'halogène ; un groupe di (alkyl en C₁-C₄) aminocarbonyl ; ou un groupe cycloalkyl (en C₁-C₄) aminocarbonyl ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsque M et R sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, ils peuvent former un cycle dans lequel MR représente la structure :


-OCH₂O-, -OCF₂O- ou



Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α-nitroacétophénone ayant la structure :



dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-di(alcoxy en C₁-C₄)éthylamine, à une température élevée, pour former un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou un α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène ayant la structure :




dans laquelle L, M, R et W sont décrits comme ci-dessus, et le traitement de l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-cyanostyrène ou de l'α-[2,2-di(alcoxy en C₁-C₄)éthylamino]-β-nitrostyrène ainsi formé avec un acide minéral ou organique pour former l'arylpyrrole désiré.

2. Procédé selon la revendication 1, dans lequel la réaction de la 2,2-di(alcoxy en C₁-C₄)éthylamine avec le benzoylacétonitrile ou l'α-nitroacétophénone est effectuée sans solvant ou en présence d'un solvant organique inerte.

3. Procédé selon la revendication 1, dans lequel on traite l' α -[2,2-di(alcoxy en C₁-C₄)éthylamino]- β -cyanostyrène ou d' α -[2,2-di(alcoxy en C₁-C₄)éthylamino]- β -nitrostyrène avec l'acide chlorhydrique, bromhydrique ou trifluoroacétique pour former l'arylpyrrole.

5 4. Procédé pour la préparation d'un α -[2,2-di(alcoxy en C₁-C₄)éthylamino]- β -cyanostyrène ou d'un α -[2,2-di(alcoxy en C₁-C₄)éthylamino]- β -nitrostyrène, représenté par la structure :




20 dans laquelle W est CN ou NO₂ ; L est H, F, Cl ou Br ; et M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsqu'ils sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, M et R peuvent former un cycle dans lequel MR représente la structure :

25 -OCH₂O-, -OCF₂O- ou




35 Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α -nitroacétophénone ayant la structure :



50 dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-diéthoxyéthylamine, à une température élevée, pour former un α -(2,2-di(alcoxy en C₁-C₄)éthylamino)- β -cyanostyrène ou un α -(2,2-di(alcoxy en C₁-C₄)éthylamino)- β -nitrostyrène ayant la structure :


55

(d)




dans laquelle L, M, R et W sont décrits comme ci-dessus.

5. Procédé pour la préparation d'un composé représenté par la structure :




dans laquelle W est CN ou NO₂, L est H, F, Cl ou Br ; M et R sont chacun indépendamment H, un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃, un groupe alkylthio en C₁-C₃, un groupe alkylsulfinyle en C₁-C₃, un groupe alkylsulfonyl en C₁-C₃, un groupe cyano, F, Cl, Br, I, un groupe nitro, CF₃, R₁CF₂Z, R₂CO ou NR₃R₄ et, lorsqu'ils sont sur des positions adjacentes, avec les atomes de carbone auxquels ils sont fixés, M et R peuvent former un cycle dans lequel MR représente la structure :


-OCH₂O-, -OCF₂O- ou



Z est S(O)_n ou O ; R₁ est H, F, CHF₂, CHFCl ou CF₃ ; R₂ est un groupe alkyle en C₁-C₃, un groupe alcoxy en C₁-C₃ ou NR₃R₄ ; R₃ est H ou un groupe alkyle en C₁-C₃ ; R₄ est H, un groupe alkyle en C₁-C₃ ou R₅CO ; R₅ est H ou un groupe alkyle en C₁-C₃ ; et n est un entier valant 0, 1 ou 2 ; comprenant la réaction d'un benzoylacétonitrile ou d'une α-nitroacétophénone ayant la structure :



dans laquelle L, M, R et W sont décrits comme ci-dessus, avec une 2,2-di(alcoxy en C₁-C₄)éthylamine, à une température élevée, pour obtenir un composé ayant la structure :



dans laquelle L, M, R et W sont décrits comme ci-dessus.